

NORTH CAROLINA

2010 School Health Profiles Report Weighted Principal Survey Results

1. Percentage of schools that ever used the School Health Index or other self-assessment tool to assess school policies, activities, and programs in the following areas.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|-----------------------------------|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Physical activity | 23.3 | 17.7 - 30.0 | 130 | 43.1 | 37.3 - 49.1 | 204 | 29.2 | 15.9 - 47.5 | 19 | 34.6 | 30.6 - 38.9 | 353 |
| b. Nutrition | 25.1 | 19.2 - 32.2 | 130 | 37.6 | 32.1 - 43.3 | 204 | 40.9 | 26.1 - 57.6 | 19 | 33.1 | 29.1 - 37.4 | 353 |
| c. Tobacco-use prevention | 29.3 | 23.2 - 36.1 | 131 | 37.6 | 32.1 - 43.4 | 202 | 46.8 | 29.3 - 65.1 | 19 | 35.1 | 31.0 - 39.4 | 352 |
| d. Asthma | 15.9 | 11.2 - 22.1 | 131 | 22.5 | 18.0 - 27.9 | 203 | 11.7 | 3.5 - 32.3 | 19 | 19.2 | 15.9 - 23.0 | 353 |
| e. Injury and violence prevention | 23.3 | 17.7 - 30.0 | 130 | 33.6 | 28.4 - 39.2 | 202 | 23.4 | 9.7 - 46.5 | 19 | 28.9 | 25.1 - 33.1 | 351 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

2. Percentage of schools with a School Improvement Plan that includes health-related goals and objectives on the following topics.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Health education | 51.9 | 46.0 - 57.8 | 131 | 55.2 | 49.4 - 60.8 | 205 | 48.2 | 31.8 - 65.0 | 19 | 53.4 | 49.3 - 57.5 | 355 |
| b. Physical education and physical activity | 50.1 | 43.9 - 56.4 | 132 | 61.0 | 55.1 - 66.7 | 205 | 48.2 | 31.0 - 65.8 | 19 | 55.9 | 51.7 - 60.1 | 356 |
| c. Nutrition services and foods and beverages available at school | 39.0 | 32.3 - 46.1 | 131 | 29.6 | 24.8 - 35.0 | 205 | 15.3 | 5.6 - 35.6 | 19 | 32.1 | 28.3 - 36.2 | 355 |
| d. Health services | 35.5 | 29.0 - 42.5 | 132 | 36.2 | 30.8 - 41.8 | 205 | 15.3 | 5.6 - 35.6 | 19 | 34.3 | 30.4 - 38.6 | 356 |
| e. Mental health and social services | 30.0 | 23.8 - 37.0 | 131 | 29.3 | 24.4 - 34.8 | 205 | 21.2 | 12.2 - 34.2 | 19 | 29.0 | 25.2 - 33.0 | 355 |
| f. Healthy and safe school environment | 83.7 | 79.1 - 87.5 | 133 | 83.9 | 79.2 - 87.7 | 206 | 73.0 | 56.5 - 84.9 | 19 | 83.0 | 79.8 - 85.8 | 358 |
| g. Family and community involvement | 79.2 | 74.1 - 83.5 | 133 | 84.4 | 80.0 - 88.0 | 206 | - | - | 18 | 81.5 | 78.3 - 84.3 | 357 |
| h. Faculty and staff health promotion | 38.9 | 32.1 - 46.0 | 132 | 38.7 | 33.5 - 44.3 | 204 | - | - | 18 | 38.3 | 34.3 - 42.6 | 354 |

- Results are suppressed due to insufficient number of respondents in subgroup.
N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

3. Percentage of schools that are required to report each of the following types of information regarding implementation of the local wellness policy.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Number of minutes of physical education required in each grade | 55.5 | 49.1 - 61.7 | 132 | 82.5 | 78.0 - 86.2 | 206 | 35.7 | 26.9 - 45.5 | 19 | 68.9 | 65.4 - 72.1 | 357 |
| b. Rates of student participation in school meal programs | 88.2 | 83.7 - 91.6 | 133 | 90.7 | 87.2 - 93.3 | 204 | 55.4 | 38.2 - 71.4 | 19 | 87.1 | 84.4 - 89.5 | 356 |
| c. Revenue from sale of foods and beverages from school-sponsored fundraisers, vending machines, school stores, or a la carte lines in the school cafeteria | 76.8 | 71.0 - 81.7 | 132 | 75.6 | 70.2 - 80.3 | 200 | 26.2 | 14.4 - 42.9 | 19 | 72.3 | 68.6 - 75.8 | 351 |
| d. Number of minutes of physical activity outside of physical education (e.g., classroom physical activity breaks, free time physical activity, or recess) | 22.9 | 18.1 - 28.6 | 132 | 64.8 | 58.9 - 70.3 | 205 | 16.7 | 7.6 - 33.0 | 19 | 45.4 | 41.6 - 49.2 | 356 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

4. Percentage of schools that currently have someone who oversees or coordinates school health and safety programs and activities.

| High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---------------------|-------------------------|-----|-----------------------|-------------------------|-----|-----------------------------------|-------------------------|----|--------------------|-------------------------|-----|
| Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| 82.3 | 75.4 - 87.5 | 132 | 87.7 | 83.6 - 90.9 | 203 | 67.1 | 55.3 - 77.1 | 19 | 84.1 | 80.7 - 87.0 | 354 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

5. Percentage of schools that have one or more than one group (e.g., a school health council, committee, or team) that offers guidance on the development of policies or coordinates activities on health topics.

| High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---------------------|-------------------------|-----|-----------------------|-------------------------|-----|-----------------------------------|-------------------------|----|--------------------|-------------------------|-----|
| Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| 47.5 | 40.1 - 54.9 | 132 | 54.0 | 48.2 - 59.7 | 205 | - | - | 18 | 49.9 | 45.4 - 54.3 | 355 |

- Results are suppressed due to insufficient number of respondents in subgroup.
N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

6. Percentage of schools that have the following groups represented on any school health council, committee, or team.*

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---|--------------|-------------------------|----|----------------|-------------------------|-----|----------------------------|-------------------------|---|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. School administrators | 82.3 | 72.7 - 89.0 | 63 | 86.6 | 80.2 - 91.2 | 111 | - | - | 6 | 84.8 | 79.6 - 88.8 | 180 |
| b. Health education teachers | 86.8 | 77.8 - 92.5 | 63 | 93.8 | 88.9 - 96.6 | 111 | - | - | 6 | 90.7 | 86.3 - 93.7 | 180 |
| c. Physical education teachers | 86.8 | 77.8 - 92.5 | 63 | 92.9 | 87.8 - 95.9 | 111 | - | - | 6 | 90.2 | 85.8 - 93.3 | 180 |
| d. Mental health or social services staff | 57.2 | 47.5 - 66.3 | 62 | 55.8 | 47.3 - 63.9 | 111 | - | - | 6 | 55.3 | 49.0 - 61.5 | 179 |
| e. Nutrition or food service staff | 51.8 | 42.3 - 61.1 | 61 | 48.6 | 41.9 - 55.3 | 111 | - | - | 6 | 49.0 | 43.5 - 54.5 | 178 |
| f. Health services staff (e.g., school nurse) | 75.2 | 64.4 - 83.5 | 62 | 82.8 | 76.2 - 87.9 | 111 | - | - | 6 | 79.4 | 73.9 - 84.0 | 179 |
| g. Maintenance and transportation staff | 19.9 | 12.6 - 29.8 | 61 | 17.8 | 12.8 - 24.3 | 111 | - | - | 6 | 17.8 | 13.6 - 22.9 | 178 |
| h. Technology staff | 24.2 | 16.1 - 34.7 | 62 | 27.7 | 21.6 - 34.8 | 111 | - | - | 6 | 26.7 | 21.7 - 32.4 | 179 |
| i. Library/media center staff | 27.3 | 19.0 - 37.6 | 62 | 26.1 | 20.1 - 33.1 | 110 | - | - | 6 | 26.0 | 21.1 - 31.5 | 178 |
| j. Student body | 50.8 | 40.7 - 60.8 | 61 | 32.8 | 25.6 - 40.8 | 110 | - | - | 6 | 40.6 | 34.7 - 46.8 | 177 |
| k. Parents or families of students | 58.7 | 48.6 - 68.1 | 62 | 56.0 | 48.3 - 63.5 | 110 | - | - | 6 | 58.0 | 52.0 - 63.8 | 178 |
| l. Community members | 51.7 | 42.2 - 61.1 | 63 | 52.8 | 45.3 - 60.2 | 111 | - | - | 6 | 52.1 | 46.3 - 57.9 | 180 |

*Among those schools that have one or more than one group that offers guidance on the development of policies or coordinates activities on health topics.

- Results are suppressed due to insufficient number of respondents in subgroup.

N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

6. Percentage of schools that have the following groups represented on any school health council, committee, or team.*

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---|--------------|-------------------------|----|----------------|-------------------------|-----|----------------------------|-------------------------|---|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| m. Local health departments, agencies, or organizations | 47.1 | 37.4 - 57.1 | 63 | 44.3 | 36.5 - 52.3 | 108 | - | - | 6 | 46.3 | 40.2 - 52.5 | 177 |
| n. Faith-based organizations | 25.2 | 16.9 - 35.7 | 62 | 15.1 | 10.5 - 21.2 | 111 | - | - | 6 | 18.0 | 13.8 - 23.2 | 179 |
| o. Businesses | 29.8 | 21.4 - 39.9 | 62 | 24.1 | 18.3 - 31.1 | 111 | - | - | 6 | 26.0 | 21.1 - 31.6 | 179 |
| p. Local government agencies | 36.4 | 27.7 - 46.1 | 63 | 31.2 | 25.0 - 38.2 | 111 | - | - | 6 | 32.3 | 27.2 - 37.9 | 180 |

*Among those schools that have one or more than one group that offers guidance on the development of policies or coordinates activities on health topics.

- Results are suppressed due to insufficient number of respondents in subgroup.

N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

7. Percentage of schools that require any school staff to receive professional development on HIV, STD, or pregnancy prevention issues and resources for the following groups.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|--|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Ethnic/racial minority youth at high risk (e.g., black, Hispanic, or American Indian youth) | 41.7 | 35.1 - 48.7 | 132 | 36.5 | 31.1 - 42.2 | 205 | 28.4 | 14.0 - 49.3 | 19 | 37.9 | 33.8 - 42.1 | 356 |
| b. Youth who participate in drop-out prevention, alternative education, or GED programs | 47.4 | 40.2 - 54.8 | 132 | 35.1 | 30.3 - 40.3 | 204 | 34.3 | 17.4 - 56.4 | 19 | 39.7 | 35.6 - 44.0 | 355 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

8. Percentage of schools that have a student-led club that aims to create a safe, welcoming, and accepting school environment for all youth, regardless of sexual orientation or gender identity.

| High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---------------------|-------------------------|-----|-----------------------|-------------------------|-----|-----------------------------------|-------------------------|----|--------------------|-------------------------|-----|
| Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| 32.4 | 26.3 - 39.1 | 133 | 11.3 | 8.0 - 15.7 | 205 | 11.7 | 3.5 - 32.3 | 19 | 19.3 | 16.2 - 22.9 | 357 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

9. Percentage of schools that engage in the following practices related to LGBTQ youth.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|--|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Identify “safe spaces” (e.g., a counselor’s office, designated classroom, or student organization) where LGBTQ youth can receive support from administrators, teachers, or other school staff | 63.2 | 55.5 - 70.3 | 133 | 44.1 | 38.5 - 49.8 | 204 | 47.4 | 32.0 - 63.3 | 19 | 51.6 | 47.3 - 55.9 | 356 |
| b. Prohibit harassment based on a student’s perceived or actual sexual orientation or gender identity | 92.1 | 86.9 - 95.3 | 133 | 80.3 | 75.3 - 84.5 | 204 | 94.2 | 74.6 - 98.9 | 19 | 85.8 | 82.5 - 88.6 | 356 |
| c. Encourage staff to attend professional development on safe and supportive school environments for all students, regardless of sexual orientation or gender identity | 54.3 | 47.7 - 60.7 | 132 | 50.1 | 44.6 - 55.7 | 205 | 64.9 | 64.9 - 64.9 | 19 | 52.8 | 48.9 - 56.7 | 356 |
| d. Facilitate access to providers not on school property who have experience in providing health services, including HIV/STD testing and counseling, to LGBTQ youth | 48.5 | 42.5 - 54.4 | 132 | 31.6 | 26.8 - 36.8 | 203 | 61.3 | 46.2 - 74.5 | 19 | 40.2 | 36.5 - 43.9 | 354 |
| e. Facilitate access to providers not on school property who have experience in providing social and psychological services to LGBTQ youth | 46.3 | 40.5 - 52.2 | 132 | 34.6 | 29.7 - 39.9 | 202 | 49.6 | 35.5 - 63.7 | 19 | 40.2 | 36.5 - 44.0 | 353 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

10. Percentage of schools that have adopted a policy that addresses each of the following issues on HIV or AIDS.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|--|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Attendance of students with HIV infection | 51.1 | 43.4 - 58.7 | 132 | 57.2 | 51.8 - 62.3 | 204 | - | - | 18 | 54.2 | 49.8 - 58.6 | 354 |
| b. Procedures to protect HIV-infected students and staff from discrimination | 65.5 | 57.9 - 72.3 | 132 | 65.2 | 59.8 - 70.2 | 204 | - | - | 18 | 65.5 | 61.2 - 69.7 | 354 |
| c. Maintaining confidentiality of HIV-infected students and staff | 76.8 | 69.6 - 82.7 | 133 | 75.1 | 70.1 - 79.6 | 205 | - | - | 18 | 76.2 | 72.2 - 79.8 | 356 |
| d. Worksite safety (i.e., universal precautions for all school staff) | 85.6 | 79.0 - 90.3 | 133 | 85.3 | 81.2 - 88.6 | 205 | 94.2 | 74.6 - 98.9 | 19 | 86.0 | 82.7 - 88.8 | 357 |
| e. Confidential counseling for HIV-infected students | 70.8 | 63.0 - 77.6 | 133 | 59.6 | 54.0 - 64.9 | 205 | - | - | 18 | 64.2 | 59.7 - 68.5 | 356 |
| f. Communication of the policy to students, school staff, and parents | 61.1 | 53.8 - 68.0 | 130 | 56.2 | 50.8 - 61.5 | 204 | - | - | 18 | 58.7 | 54.3 - 62.9 | 352 |
| g. Adequate training about HIV infection for school staff | 65.2 | 57.2 - 72.5 | 132 | 59.0 | 53.3 - 64.6 | 205 | - | - | 18 | 62.4 | 57.9 - 66.7 | 355 |
| h. Procedures for implementing the policy | 59.8 | 52.1 - 67.0 | 132 | 54.3 | 48.6 - 60.0 | 204 | - | - | 18 | 56.7 | 52.1 - 61.2 | 354 |

- Results are suppressed due to insufficient number of respondents in subgroup.

N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

11. Percentage of schools that have or participate in each of the following programs.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. A student mentoring program | 75.6 | 69.0 - 81.2 | 133 | 71.1 | 66.3 - 75.5 | 205 | 84.7 | 77.9 - 89.6 | 19 | 73.8 | 70.2 - 77.2 | 357 |
| b. A safe-passages to school program | 18.9 | 14.0 - 25.1 | 132 | 21.1 | 16.7 - 26.3 | 204 | 27.0 | 15.1 - 43.5 | 19 | 20.7 | 17.4 - 24.4 | 355 |
| c. A program to prevent bullying | 68.7 | 61.7 - 75.0 | 133 | 92.2 | 88.7 - 94.7 | 202 | 88.3 | 67.7 - 96.5 | 19 | 82.9 | 79.5 - 85.9 | 354 |
| d. A program to prevent dating violence | 35.0 | 28.3 - 42.3 | 130 | 19.5 | 15.5 - 24.4 | 205 | 46.0 | 27.2 - 66.0 | 19 | 27.3 | 23.6 - 31.4 | 354 |
| e. A youth development program | 48.3 | 41.4 - 55.3 | 131 | 46.9 | 41.7 - 52.2 | 205 | 53.2 | 40.1 - 66.0 | 19 | 47.9 | 43.9 - 51.9 | 355 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

12. Percentage of schools in which all staff who teach health education topics are certified, licensed, or endorsed by the state in health education.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|-------------------|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Yes | 87.4 | 82.7 - 91.0 | 132 | 84.7 | 80.3 - 88.2 | 204 | 73.0 | 56.5 - 84.9 | 19 | 84.9 | 81.8 - 87.5 | 355 |
| b. No | 8.4 | 4.9 - 14.0 | 132 | 12.8 | 9.7 - 16.8 | 204 | 21.2 | 12.2 - 34.2 | 19 | 11.8 | 9.3 - 14.7 | 355 |
| c. Not applicable | 4.2 | 1.8 - 9.2 | 132 | 2.5 | 1.2 - 5.1 | 204 | 5.8 | 1.1 - 25.4 | 19 | 3.4 | 2.0 - 5.6 | 355 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

13. Percentage of schools that require physical education for students in any of grades 6 through 12.

| High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---------------------|-------------------------------|-----|-----------------------|-------------------------------|-----|-----------------------------------|-------------------------------|----|--------------------|-------------------------------|-----|
| Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| 97.2 | 93.8 - 98.7 | 133 | 98.1 | 95.7 - 99.1 | 205 | 88.3 | 67.7 - 96.5 | 19 | 97.0 | 95.0 - 98.2 | 357 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

14. Percentage of schools that taught a required physical education course in each of the following grades.*

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|-------------------|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Sixth grade | § | § | § | 96.4 | 93.5 - 98.1 | 195 | 87.4 | 65.4 - 96.2 | 17 | 95.4 | 92.0 - 97.3 | 212 |
| b. Seventh grade | § | § | § | 95.6 | 92.5 - 97.4 | 204 | 87.9 | 66.6 - 96.3 | 18 | 94.7 | 91.4 - 96.8 | 222 |
| c. Eighth grade | § | § | § | 95.6 | 92.5 - 97.4 | 204 | 87.9 | 66.6 - 96.3 | 18 | 94.7 | 91.4 - 96.8 | 222 |
| d. Ninth grade | 94.5 | 89.9 - 97.1 | 130 | 20.2 | 8.3 - 41.6 | 15 | 76.6 | 61.0 - 87.3 | 19 | 85.7 | 80.9 - 89.4 | 164 |
| e. Tenth grade | 11.4 | 7.5 - 17.0 | 125 | § | § | § | 30.6 | 14.4 - 53.7 | 19 | 14.7 | 10.2 - 20.7 | 144 |
| f. Eleventh grade | 9.2 | 5.5 - 14.8 | 125 | § | § | § | 6.7 | 1.3 - 28.8 | 16 | 8.8 | 5.4 - 14.0 | 141 |
| g. Twelfth grade | 9.2 | 5.5 - 14.8 | 125 | § | § | § | 6.7 | 1.3 - 28.8 | 16 | 8.8 | 5.4 - 14.0 | 141 |

*Among schools with students in that grade.

§ Grade is not applicable for type of school.

N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

15. Percentage of schools in which students can be exempted from taking required physical education for one grading period or longer for any of the following reasons.*

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Enrollment in other courses (e.g., math or science) | 24.7 | 19.1 - 31.1 | 128 | 12.3 | 8.9 - 16.8 | 201 | 26.5 | 14.0 - 44.4 | 17 | 17.9 | 14.9 - 21.5 | 346 |
| b. Participation in school sports | 3.4 | 1.5 - 7.6 | 128 | 2.0 | 0.9 - 4.5 | 201 | 6.6 | 1.2 - 28.4 | 17 | 2.8 | 1.6 - 4.9 | 346 |
| c. Participation in other school activities (e.g., ROTC, band, or chorus) | 11.1 | 7.3 - 16.6 | 128 | 10.8 | 7.6 - 15.1 | 201 | 13.2 | 11.4 - 15.4 | 17 | 11.1 | 8.7 - 14.1 | 346 |
| d. Participation in community sports activities | 1.9 | 0.6 - 6.0 | 128 | 1.5 | 0.6 - 3.7 | 201 | 0.0 | † | 17 | 1.5 | 0.7 - 3.2 | 346 |
| e. Religious reasons | 28.6 | 22.6 - 35.4 | 126 | 38.6 | 33.2 - 44.2 | 199 | 6.6 | 1.2 - 28.4 | 17 | 32.6 | 28.8 - 36.7 | 342 |
| f. Long-term physical or medical disability | 60.4 | 52.9 - 67.5 | 128 | 74.6 | 69.3 - 79.2 | 199 | 49.5 | 39.7 - 59.4 | 17 | 67.5 | 63.4 - 71.3 | 344 |
| g. Cognitive disability | 30.1 | 23.8 - 37.2 | 128 | 21.3 | 17.2 - 26.1 | 200 | 4.1 | 0.8 - 18.7 | 17 | 23.5 | 20.1 - 27.2 | 345 |
| h. High physical fitness competency test score | 1.5 | 0.5 - 4.5 | 128 | 0.5 | 0.1 - 2.3 | 201 | 0.0 | † | 17 | 0.8 | 0.3 - 2.1 | 346 |
| i. Participation in vocational training | 3.7 | 1.8 - 7.4 | 128 | 2.4 | 1.2 - 4.9 | 201 | 0.0 | † | 17 | 2.7 | 1.6 - 4.5 | 346 |
| j. Participation in community service activities | 1.5 | 0.5 - 4.5 | 128 | 1.0 | 0.3 - 3.0 | 201 | 0.0 | † | 17 | 1.1 | 0.5 - 2.4 | 346 |

*Among those schools that require physical education for students in any of grades 6 through 12.

† Confidence intervals not applicable for 0 percent or 100 percent.

N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

16. Percentage of schools in which physical education teachers or specialists received professional development on physical education during the past two years.

| High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---------------------|-------------------------------|-----|-----------------------|-------------------------------|-----|-----------------------------------|-------------------------------|----|--------------------|-------------------------------|-----|
| Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| 90.1 | 86.0 - 93.1 | 130 | 95.5 | 93.3 - 97.0 | 205 | - | - | 17 | 92.0 | 90.2 - 93.5 | 352 |

- Results are suppressed due to insufficient number of respondents in subgroup.
N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

17. Percentage of schools that provide those who teach physical education with the following materials.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Goals, objectives, and expected outcomes for physical education | 97.0 | 95.6 - 97.9 | 132 | 98.5 | 96.3 - 99.4 | 205 | 82.5 | 71.1 - 90.0 | 19 | 96.7 | 95.4 - 97.7 | 356 |
| b. A chart describing the annual scope and sequence of instruction for physical education | 82.2 | 75.8 - 87.2 | 132 | 84.9 | 80.7 - 88.3 | 204 | 67.1 | 48.3 - 81.7 | 19 | 82.5 | 79.1 - 85.5 | 355 |
| c. Plans for how to assess student performance in physical education | 85.5 | 80.1 - 89.5 | 132 | 84.9 | 80.8 - 88.3 | 204 | 73.0 | 50.0 - 88.0 | 19 | 84.2 | 80.9 - 87.1 | 355 |
| d. A written physical education curriculum | 94.0 | 90.2 - 96.4 | 133 | 93.1 | 89.8 - 95.4 | 204 | 73.0 | 56.5 - 84.9 | 19 | 92.0 | 89.5 - 93.9 | 356 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

18. Percentage of schools that offer opportunities for all students to participate in intramural activities or physical activity clubs.

| High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---------------------|-------------------------|-----|-----------------------|-------------------------|-----|-----------------------------------|-------------------------|----|--------------------|-------------------------|-----|
| Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| 51.9 | 45.2 - 58.6 | 133 | 67.1 | 61.5 - 72.3 | 204 | 57.7 | 37.1 - 75.9 | 19 | 60.6 | 56.4 - 64.8 | 356 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

19. Percentage of schools in which children or adolescents use indoor physical activity or athletic facilities for community-sponsored physical activity classes or lessons.

| High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---------------------|-------------------------|-----|-----------------------|-------------------------|-----|-----------------------------------|-------------------------|----|--------------------|-------------------------|-----|
| Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| 69.9 | 62.9 - 76.0 | 131 | 75.1 | 70.0 - 79.5 | 205 | 48.2 | 28.0 - 69.0 | 19 | 71.1 | 67.0 - 74.9 | 355 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

20. Percentage of schools that have adopted a policy prohibiting tobacco use.

| High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---------------------|-------------------------------|-----|-----------------------|-------------------------------|-----|-----------------------------------|-------------------------------|----|--------------------|-------------------------------|-----|
| Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| 99.3 | 96.6 - 99.9 | 133 | 99.0 | 96.9 - 99.7 | 206 | 94.2 | 74.6 - 98.9 | 19 | 98.7 | 97.2 - 99.5 | 358 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

21. (Students) Percentage of schools that have a tobacco-use prevention policy that specifically prohibits the use of each type of tobacco for students during any school-related activity.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|----------------------|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Cigarettes | 98.6 | 95.6 - 99.5 | 132 | 99.0 | 96.9 - 99.7 | 206 | 94.2 | 74.6 - 98.9 | 19 | 98.5 | 96.9 - 99.3 | 357 |
| b. Smokeless Tobacco | 98.6 | 95.6 - 99.5 | 132 | 98.5 | 96.2 - 99.4 | 206 | 94.2 | 74.6 - 98.9 | 19 | 98.2 | 96.5 - 99.1 | 357 |
| c. Cigars | 98.6 | 95.6 - 99.5 | 132 | 98.0 | 95.6 - 99.1 | 206 | 94.2 | 74.6 - 98.9 | 19 | 97.9 | 96.2 - 98.9 | 357 |
| d. Pipes | 98.6 | 95.6 - 99.5 | 132 | 98.0 | 95.6 - 99.1 | 206 | 94.2 | 74.6 - 98.9 | 19 | 97.9 | 96.2 - 98.9 | 357 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

21. (Faculty/Staff) Percentage of schools that have a tobacco-use prevention policy that specifically prohibits the use of each type of tobacco for faculty/staff during any school-related activity.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|----------------------|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Cigarettes | 98.6 | 95.6 - 99.5 | 132 | 99.0 | 96.9 - 99.7 | 204 | 94.2 | 74.6 - 98.9 | 19 | 98.5 | 96.8 - 99.3 | 355 |
| b. Smokeless Tobacco | 98.6 | 95.6 - 99.5 | 132 | 98.5 | 96.2 - 99.4 | 204 | 94.2 | 74.6 - 98.9 | 19 | 98.2 | 96.5 - 99.1 | 355 |
| c. Cigars | 98.6 | 95.6 - 99.5 | 132 | 98.0 | 95.5 - 99.1 | 204 | 94.2 | 74.6 - 98.9 | 19 | 97.9 | 96.2 - 98.9 | 355 |
| d. Pipes | 98.6 | 95.6 - 99.5 | 132 | 98.0 | 95.5 - 99.1 | 204 | 94.2 | 74.6 - 98.9 | 19 | 97.9 | 96.2 - 98.9 | 355 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

21. (Visitors) Percentage of schools that have a tobacco-use prevention policy that specifically prohibits the use of each type of tobacco for visitors during any school-related activity.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|----------------------|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Cigarettes | 98.6 | 95.6 - 99.5 | 131 | 98.5 | 96.2 - 99.4 | 203 | 94.2 | 74.6 - 98.9 | 19 | 98.2 | 96.5 - 99.1 | 353 |
| b. Smokeless Tobacco | 98.6 | 95.6 - 99.5 | 131 | 98.0 | 95.5 - 99.1 | 202 | 94.2 | 74.6 - 98.9 | 19 | 97.9 | 96.1 - 98.9 | 352 |
| c. Cigars | 98.6 | 95.6 - 99.5 | 131 | 97.4 | 94.8 - 98.7 | 200 | 94.2 | 74.6 - 98.9 | 19 | 97.6 | 95.8 - 98.7 | 350 |
| d. Pipes | 98.6 | 95.6 - 99.5 | 131 | 97.4 | 94.8 - 98.8 | 201 | 94.2 | 74.6 - 98.9 | 19 | 97.6 | 95.8 - 98.7 | 351 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

22. (Students) Percentage of schools that have a tobacco-use prevention policy that specifically prohibits tobacco use during each of the following times for students.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|----------------------------|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. During school hours | 99.3 | 96.5 - 99.9 | 131 | 98.0 | 95.6 - 99.1 | 206 | 94.2 | 74.6 - 98.9 | 19 | 98.2 | 96.5 - 99.1 | 356 |
| b. During non-school hours | 96.7 | 92.6 - 98.6 | 131 | 93.7 | 90.5 - 95.8 | 206 | 88.3 | 67.7 - 96.5 | 19 | 94.4 | 92.0 - 96.1 | 356 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

22. (Faculty/Staff) Percentage of schools that have a tobacco-use prevention policy that specifically prohibits tobacco use during each of the following times for faculty/staff.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|----------------------------|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. During school hours | 99.3 | 96.5 - 99.9 | 131 | 98.0 | 95.6 - 99.1 | 206 | 94.2 | 74.6 - 98.9 | 19 | 98.2 | 96.5 - 99.1 | 356 |
| b. During non-school hours | 96.0 | 91.7 - 98.1 | 131 | 91.7 | 88.1 - 94.3 | 206 | 88.3 | 67.7 - 96.5 | 19 | 93.0 | 90.5 - 95.0 | 356 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

22. (Visitors) Percentage of schools that have a tobacco-use prevention policy that specifically prohibits tobacco use during each of the following times for visitors.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|----------------------------|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. During school hours | 99.3 | 96.5 - 99.9 | 131 | 97.5 | 94.9 - 98.8 | 206 | 94.2 | 74.6 - 98.9 | 19 | 97.9 | 96.2 - 98.9 | 356 |
| b. During non-school hours | 96.0 | 91.7 - 98.1 | 131 | 91.2 | 87.5 - 93.9 | 206 | 88.3 | 67.7 - 96.5 | 19 | 92.8 | 90.1 - 94.7 | 356 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

23. (Students) Percentage of schools that have a tobacco-use prevention policy that specifically prohibits tobacco use in each of the following locations for students.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. In school buildings | 99.3 | 96.5 - 99.9 | 132 | 99.0 | 96.9 - 99.7 | 206 | 94.2 | 74.6 - 98.9 | 19 | 98.7 | 97.2 - 99.4 | 357 |
| b. Outside on school grounds, including parking lots and playing fields | 99.3 | 96.5 - 99.9 | 132 | 99.0 | 96.9 - 99.7 | 206 | 94.2 | 74.6 - 98.9 | 19 | 98.7 | 97.2 - 99.4 | 357 |
| c. On school buses or other vehicles used to transport students | 99.3 | 96.5 - 99.9 | 132 | 98.5 | 96.2 - 99.4 | 206 | 94.2 | 74.6 - 98.9 | 19 | 98.5 | 96.8 - 99.3 | 357 |
| d. At off-campus, school-sponsored events | 97.4 | 93.4 - 99.0 | 131 | 96.0 | 92.9 - 97.7 | 204 | 88.3 | 67.7 - 96.5 | 19 | 95.9 | 93.6 - 97.4 | 354 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

23. (Faculty/Staff) Percentage of schools that have a tobacco-use prevention policy that specifically prohibits tobacco use in each of the following locations for faculty/staff.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. In school buildings | 99.3 | 96.5 - 99.9 | 131 | 99.0 | 96.9 - 99.7 | 206 | 94.2 | 74.6 - 98.9 | 19 | 98.7 | 97.2 - 99.4 | 356 |
| b. Outside on school grounds, including parking lots and playing fields | 99.3 | 96.5 - 99.9 | 131 | 99.0 | 96.9 - 99.7 | 205 | 94.2 | 74.6 - 98.9 | 19 | 98.7 | 97.2 - 99.4 | 355 |
| c. On school buses or other vehicles used to transport students | 99.3 | 96.5 - 99.9 | 131 | 98.5 | 96.2 - 99.4 | 206 | 94.2 | 74.6 - 98.9 | 19 | 98.5 | 96.8 - 99.3 | 356 |
| d. At off-campus, school-sponsored events | 95.2 | 90.8 - 97.6 | 130 | 92.1 | 88.6 - 94.6 | 204 | 88.3 | 67.7 - 96.5 | 19 | 93.0 | 90.4 - 94.9 | 353 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

23. (Visitors) Percentage of schools that have a tobacco-use prevention policy that specifically prohibits tobacco use in each of the following locations for visitors.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. In school buildings | 99.3 | 96.5 - 99.9 | 130 | 98.5 | 96.2 - 99.4 | 203 | 94.2 | 74.6 - 98.9 | 19 | 98.5 | 96.8 - 99.3 | 352 |
| b. Outside on school grounds, including parking lots and playing fields | 99.3 | 96.5 - 99.9 | 129 | 98.5 | 96.2 - 99.4 | 203 | 94.2 | 74.6 - 98.9 | 19 | 98.4 | 96.8 - 99.3 | 351 |
| c. On school buses or other vehicles used to transport students | 99.3 | 96.5 - 99.9 | 130 | 98.0 | 95.5 - 99.1 | 203 | 93.8 | 73.2 - 98.8 | 18 | 98.2 | 96.4 - 99.1 | 351 |
| d. At off-campus, school-sponsored events | 82.4 | 76.2 - 87.2 | 129 | 82.5 | 78.1 - 86.2 | 201 | 76.6 | 61.0 - 87.3 | 19 | 82.0 | 78.6 - 85.0 | 349 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

23N. The percentage of schools that follow a policy that mandates a “tobacco-free environment.” A “tobacco-free environment” is one that prohibits tobacco use by students, staff, and visitors in school buildings, at school functions, in school vehicles, on school grounds, and at off-site school events, applicable 24 hours a day and seven days a week.*

| High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---------------------|-------------------------|-----|-----------------------|-------------------------|-----|-----------------------------------|-------------------------|----|--------------------|-------------------------|-----|
| Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| 77.9 | 70.9 - 83.6 | 127 | 77.1 | 71.9 - 81.5 | 197 | 70.8 | 60.6 - 79.2 | 19 | 76.9 | 73.1 - 80.3 | 343 |

*Responses to question 21 (a, b, c, and d), question 22 (a and b), and question 23 (a, b, c, and d) are all "yes."
 N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

24. Percentage of schools that have procedures to inform each of the following groups about the tobacco-use prevention policy that prohibits their use of tobacco.*

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|----------------------|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Students | 100.0 | † | 131 | 98.5 | 96.2 - 99.4 | 204 | 100.0 | † | 18 | 99.2 | 97.9 - 99.7 | 353 |
| b. Faculty and staff | 100.0 | † | 131 | 98.5 | 96.3 - 99.4 | 204 | 100.0 | † | 18 | 99.2 | 98.0 - 99.7 | 353 |
| c. Visitors | 99.3 | 96.4 - 99.9 | 128 | 97.5 | 95.0 - 98.8 | 202 | 100.0 | † | 18 | 98.4 | 96.9 - 99.2 | 348 |

*Among those schools that have adopted a policy prohibiting tobacco use.

† Confidence intervals not applicable for 0 percent or 100 percent.

N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

25. Percentage of schools that have a tobacco-use prevention policy that includes guidelines on what actions the school should take when students are caught smoking cigarettes.*

| High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---------------------|-------------------------------|-----|-----------------------|-------------------------------|-----|-----------------------------------|-------------------------------|----|--------------------|-------------------------------|-----|
| Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| 98.1 | 94.1 - 99.4 | 131 | 97.5 | 95.0 - 98.8 | 203 | 100.0 | † | 18 | 97.9 | 96.2 - 98.9 | 352 |

*Among those schools that have adopted a policy prohibiting tobacco use.

† Confidence intervals not applicable for 0 percent or 100 percent.

N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

26. Percentage of schools in which the following person is responsible for enforcing the tobacco-use prevention policy.*

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. No single individual is responsible | 56.4 | 49.1 - 63.5 | 122 | 47.4 | 41.6 - 53.3 | 194 | 58.9 | 34.9 - 79.3 | 18 | 51.6 | 47.1 - 56.2 | 334 |
| b. Principal | 30.1 | 23.6 - 37.4 | 122 | 45.8 | 40.1 - 51.6 | 194 | 34.9 | 17.0 - 58.4 | 18 | 39.1 | 34.8 - 43.6 | 334 |
| c. Assistant principal | 13.5 | 9.3 - 19.2 | 122 | 4.1 | 2.3 - 7.2 | 194 | 6.2 | 1.2 - 26.8 | 18 | 7.8 | 5.7 - 10.5 | 334 |
| d. Other school administrator | 0.0 | † | 122 | 1.6 | 0.6 - 4.0 | 194 | 0.0 | † | 18 | 0.9 | 0.4 - 2.2 | 334 |
| e. Other school faculty or staff member | 0.0 | † | 122 | 1.0 | 0.3 - 3.2 | 194 | 0.0 | † | 18 | 0.6 | 0.2 - 1.8 | 334 |

*Among those schools that have adopted a policy prohibiting tobacco use.

† Confidence intervals not applicable for 0 percent or 100 percent.

N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

27. Percentage of schools in which the following criteria help determine what actions are taken when students are caught smoking cigarettes.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|--|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Zero tolerance | 86.1 | 80.2 - 90.5 | 133 | 87.1 | 82.8 - 90.5 | 204 | - | - | 18 | 85.4 | 82.0 - 88.2 | 355 |
| b. Effect or severity of the violation | 55.9 | 48.9 - 62.8 | 130 | 61.5 | 55.7 - 66.9 | 200 | - | - | 18 | 60.5 | 56.1 - 64.7 | 348 |
| c. Grade level of student | 20.5 | 15.1 - 27.3 | 130 | 38.8 | 32.9 - 45.1 | 199 | - | - | 18 | 33.3 | 29.0 - 37.8 | 347 |
| d. Repeat offender status | 86.1 | 80.6 - 90.3 | 130 | 78.3 | 73.1 - 82.7 | 200 | - | - | 18 | 80.8 | 77.0 - 84.0 | 348 |

- Results are suppressed due to insufficient number of respondents in subgroup.
N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

28. (Never) Percentage of schools that NEVER take the following actions when students are caught smoking cigarettes.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Parents or guardians are notified | 0.0 | † | 133 | 1.5 | 0.6 - 3.7 | 206 | - | - | 18 | 1.3 | 0.5 - 2.8 | 357 |
| b. Referred to a school counselor | 9.3 | 6.0 - 14.3 | 133 | 4.5 | 2.6 - 7.6 | 206 | - | - | 18 | 7.3 | 5.3 - 10.0 | 357 |
| c. Referred to a school administrator | 0.0 | † | 133 | 1.5 | 0.6 - 3.7 | 205 | - | - | 18 | 1.7 | 0.8 - 3.5 | 356 |
| d. Encouraged, but not required, to participate in an assistance, education, or cessation program | 25.9 | 20.8 - 31.8 | 132 | 23.1 | 18.4 - 28.6 | 204 | - | - | 18 | 24.6 | 21.0 - 28.5 | 354 |
| e. Required to participate in an assistance, education, or cessation program | 36.4 | 29.5 - 43.9 | 133 | 36.9 | 32.2 - 42.0 | 204 | - | - | 18 | 36.9 | 32.8 - 41.1 | 355 |
| f. Referred to legal authorities | 53.6 | 46.8 - 60.3 | 133 | 45.9 | 40.5 - 51.4 | 204 | - | - | 18 | 48.1 | 44.0 - 52.2 | 355 |
| g. Placed in detention | 22.5 | 17.9 - 27.9 | 133 | 25.8 | 21.0 - 31.3 | 203 | - | - | 18 | 24.5 | 21.0 - 28.3 | 354 |
| h. Not allowed to participate in extra-curricular activities or interscholastic sports | 31.8 | 25.9 - 38.4 | 132 | 24.1 | 19.6 - 29.2 | 205 | - | - | 18 | 27.8 | 24.1 - 31.8 | 355 |
| i. Given in-school suspension | 18.0 | 12.8 - 24.7 | 130 | 14.9 | 11.5 - 19.0 | 204 | - | - | 18 | 17.0 | 13.9 - 20.6 | 352 |
| j. Suspended from school | 6.4 | 4.0 - 10.2 | 132 | 8.5 | 5.7 - 12.4 | 202 | - | - | 18 | 8.0 | 5.9 - 10.7 | 352 |
| k. Expelled from school | 81.3 | 74.8 - 86.5 | 132 | 77.6 | 72.4 - 82.0 | 205 | - | - | 18 | 78.0 | 74.3 - 81.2 | 355 |
| l. Reassigned to an alternative school | 61.1 | 53.7 - 68.0 | 132 | 68.4 | 63.2 - 73.1 | 205 | - | - | 18 | 65.0 | 60.8 - 68.8 | 355 |

- Results are suppressed due to insufficient number of respondents in subgroup.

† Confidence intervals not applicable for 0 percent or 100 percent.

N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

28. (Rarely) Percentage of schools that RARELY take the following actions when students are caught smoking cigarettes.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Parents or guardians are notified | 0.0 | † | 133 | 0.0 | † | 206 | - | - | 18 | 0.4 | 0.1 - 2.1 | 357 |
| b. Referred to a school counselor | 29.0 | 22.7 - 36.2 | 133 | 10.6 | 7.5 - 14.8 | 206 | - | - | 18 | 18.1 | 14.9 - 21.9 | 357 |
| c. Referred to a school administrator | 0.0 | † | 133 | 0.0 | † | 205 | - | - | 18 | 0.0 | † | 356 |
| d. Encouraged, but not required, to participate in an assistance, education, or cessation program | 18.8 | 13.9 - 24.9 | 132 | 16.6 | 12.7 - 21.2 | 204 | - | - | 18 | 17.1 | 14.1 - 20.7 | 354 |
| e. Required to participate in an assistance, education, or cessation program | 23.2 | 17.9 - 29.4 | 133 | 19.0 | 14.9 - 24.0 | 204 | - | - | 18 | 20.2 | 16.9 - 23.9 | 355 |
| f. Referred to legal authorities | 26.3 | 20.3 - 33.3 | 133 | 31.0 | 26.3 - 36.2 | 204 | - | - | 18 | 30.0 | 26.3 - 34.1 | 355 |
| g. Placed in detention | 9.6 | 6.3 - 14.5 | 133 | 15.7 | 12.2 - 20.0 | 203 | - | - | 18 | 12.8 | 10.4 - 15.7 | 354 |
| h. Not allowed to participate in extra-curricular activities or interscholastic sports | 24.5 | 19.3 - 30.5 | 132 | 24.3 | 20.2 - 28.9 | 205 | - | - | 18 | 23.4 | 20.2 - 26.8 | 355 |
| i. Given in-school suspension | 8.9 | 5.3 - 14.3 | 130 | 10.2 | 7.3 - 14.2 | 204 | - | - | 18 | 9.5 | 7.3 - 12.4 | 352 |
| j. Suspended from school | 16.6 | 12.0 - 22.6 | 132 | 16.1 | 12.3 - 20.8 | 202 | - | - | 18 | 15.7 | 12.8 - 19.1 | 352 |
| k. Expelled from school | 12.3 | 8.1 - 18.2 | 132 | 14.1 | 10.6 - 18.7 | 205 | - | - | 18 | 13.7 | 11.0 - 17.0 | 355 |
| l. Reassigned to an alternative school | 23.5 | 17.7 - 30.6 | 132 | 21.3 | 17.1 - 26.1 | 205 | - | - | 18 | 21.9 | 18.5 - 25.9 | 355 |

- Results are suppressed due to insufficient number of respondents in subgroup.

† Confidence intervals not applicable for 0 percent or 100 percent.

N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

28. (Sometimes) Percentage of schools that SOMETIMES take the following actions when students are caught smoking cigarettes.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Parents or guardians are notified | 9.1 | 5.5 - 14.7 | 133 | 0.0 | † | 206 | - | - | 18 | 4.3 | 2.9 - 6.4 | 357 |
| b. Referred to a school counselor | 42.0 | 35.3 - 49.0 | 133 | 50.4 | 44.9 - 55.9 | 206 | - | - | 18 | 47.2 | 42.9 - 51.5 | 357 |
| c. Referred to a school administrator | 2.1 | 0.8 - 5.3 | 133 | 1.9 | 0.9 - 4.3 | 205 | - | - | 18 | 3.2 | 2.1 - 4.8 | 356 |
| d. Encouraged, but not required, to participate in an assistance, education, or cessation program | 34.2 | 27.9 - 41.1 | 132 | 28.0 | 23.0 - 33.5 | 204 | - | - | 18 | 30.8 | 26.8 - 35.1 | 354 |
| e. Required to participate in an assistance, education, or cessation program | 18.8 | 14.2 - 24.5 | 133 | 23.3 | 18.7 - 28.8 | 204 | - | - | 18 | 21.5 | 18.1 - 25.4 | 355 |
| f. Referred to legal authorities | 15.9 | 11.0 - 22.3 | 133 | 17.6 | 13.8 - 22.2 | 204 | - | - | 18 | 16.8 | 13.8 - 20.4 | 355 |
| g. Placed in detention | 40.7 | 34.4 - 47.4 | 133 | 38.4 | 33.0 - 44.2 | 203 | - | - | 18 | 40.4 | 36.2 - 44.7 | 354 |
| h. Not allowed to participate in extra-curricular activities or interscholastic sports | 34.5 | 27.8 - 41.9 | 132 | 38.0 | 32.6 - 43.8 | 205 | - | - | 18 | 36.7 | 32.4 - 41.3 | 355 |
| i. Given in-school suspension | 47.4 | 40.1 - 54.9 | 130 | 50.9 | 45.4 - 56.3 | 204 | - | - | 18 | 49.6 | 45.2 - 54.0 | 352 |
| j. Suspended from school | 51.2 | 45.0 - 57.4 | 132 | 53.2 | 47.4 - 58.9 | 202 | - | - | 18 | 51.6 | 47.3 - 55.8 | 352 |
| k. Expelled from school | 5.0 | 2.7 - 9.0 | 132 | 6.4 | 4.1 - 9.8 | 205 | - | - | 18 | 6.3 | 4.4 - 8.8 | 355 |
| l. Reassigned to an alternative school | 14.0 | 9.7 - 19.6 | 132 | 9.4 | 6.7 - 13.0 | 205 | - | - | 18 | 11.6 | 9.1 - 14.6 | 355 |

- Results are suppressed due to insufficient number of respondents in subgroup.

† Confidence intervals not applicable for 0 percent or 100 percent.

N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

28. (Always) Percentage of schools that ALWAYS or ALMOST ALWAYS take the following actions when students are caught smoking cigarettes.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Parents or guardians are notified | 90.9 | 85.3 - 94.5 | 133 | 98.5 | 96.3 - 99.4 | 206 | - | - | 18 | 94.0 | 91.5 - 95.8 | 357 |
| b. Referred to a school counselor | 19.7 | 14.1 - 26.7 | 133 | 34.6 | 29.2 - 40.3 | 206 | - | - | 18 | 27.4 | 23.5 - 31.6 | 357 |
| c. Referred to a school administrator | 97.9 | 94.7 - 99.2 | 133 | 96.6 | 93.7 - 98.1 | 205 | - | - | 18 | 95.1 | 93.0 - 96.7 | 356 |
| d. Encouraged, but not required, to participate in an assistance, education, or cessation program | 21.1 | 16.0 - 27.3 | 132 | 32.4 | 27.3 - 37.9 | 204 | - | - | 18 | 27.5 | 23.8 - 31.5 | 354 |
| e. Required to participate in an assistance, education, or cessation program | 21.6 | 16.4 - 27.9 | 133 | 20.7 | 16.3 - 25.8 | 204 | - | - | 18 | 21.4 | 18.0 - 25.3 | 355 |
| f. Referred to legal authorities | 4.2 | 2.2 - 8.1 | 133 | 5.4 | 3.3 - 8.7 | 204 | - | - | 18 | 5.0 | 3.4 - 7.3 | 355 |
| g. Placed in detention | 27.2 | 21.6 - 33.6 | 133 | 20.0 | 15.7 - 25.2 | 203 | - | - | 18 | 22.3 | 18.9 - 26.1 | 354 |
| h. Not allowed to participate in extra-curricular activities or interscholastic sports | 9.2 | 6.1 - 13.8 | 132 | 13.6 | 10.2 - 17.9 | 205 | - | - | 18 | 12.1 | 9.6 - 15.2 | 355 |
| i. Given in-school suspension | 25.7 | 19.8 - 32.7 | 130 | 24.0 | 19.9 - 28.7 | 204 | - | - | 18 | 23.8 | 20.4 - 27.6 | 352 |
| j. Suspended from school | 25.8 | 20.2 - 32.3 | 132 | 22.2 | 17.5 - 27.7 | 202 | - | - | 18 | 24.7 | 21.1 - 28.8 | 352 |
| k. Expelled from school | 1.4 | 0.5 - 4.4 | 132 | 1.9 | 0.9 - 4.3 | 205 | - | - | 18 | 2.0 | 1.1 - 3.8 | 355 |
| l. Reassigned to an alternative school | 1.4 | 0.5 - 4.4 | 132 | 1.0 | 0.3 - 3.0 | 205 | - | - | 18 | 1.5 | 0.7 - 3.1 | 355 |

- Results are suppressed due to insufficient number of respondents in subgroup.
N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

29. Percentage of schools that post signs marking a tobacco-free school zone, that is, a specified distance from school grounds where tobacco use is not allowed.

| High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---------------------|-------------------------|-----|-----------------------|-------------------------|-----|-----------------------------------|-------------------------|----|--------------------|-------------------------|-----|
| Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| 94.2 | 91.1 - 96.3 | 133 | 90.8 | 87.0 - 93.5 | 206 | 96.4 | 83.4 - 99.3 | 19 | 92.5 | 90.2 - 94.3 | 358 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

30. Percentage of schools that did the following activities during the past two years.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Gathered and shared information with students and families about mass-media messages or community-based tobacco-use prevention efforts | 47.6 | 40.8 - 54.6 | 133 | 52.9 | 46.9 - 58.8 | 206 | 42.3 | 27.9 - 58.2 | 19 | 50.1 | 45.8 - 54.5 | 358 |
| b. Worked with local agencies or organizations to plan and implement events or programs intended to reduce tobacco use | 59.7 | 52.5 - 66.4 | 133 | 49.6 | 44.0 - 55.2 | 206 | 51.8 | 34.2 - 69.0 | 19 | 53.6 | 49.3 - 57.8 | 358 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

31. Percentage of schools that provide tobacco cessation services for each of the following groups.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|----------------------|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Faculty and staff | 32.7 | 25.8 - 40.4 | 133 | 33.0 | 27.9 - 38.4 | 206 | 36.5 | 20.8 - 55.8 | 19 | 33.1 | 29.0 - 37.5 | 358 |
| b. Students | 42.8 | 36.4 - 49.5 | 133 | 35.5 | 30.5 - 40.9 | 206 | 24.8 | 11.1 - 46.7 | 19 | 37.5 | 33.6 - 41.6 | 358 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

32. Percentage of schools that have arrangements with any organizations or health care professionals not on school property to provide tobacco cessation services for each of the following groups.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|----------------------|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Faculty and staff | 43.7 | 37.2 - 50.3 | 133 | 47.4 | 42.0 - 52.9 | 205 | - | - | 18 | 46.5 | 42.4 - 50.7 | 356 |
| b. Students | 50.0 | 43.4 - 56.7 | 133 | 44.9 | 39.9 - 50.1 | 204 | - | - | 17 | 46.7 | 42.5 - 50.9 | 354 |

- Results are suppressed due to insufficient number of respondents in subgroup.
N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

33. Percentage of schools that never, rarely, sometimes, or always or almost always offer fruits or non-fried vegetables at school celebrations when foods or beverages are offered.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|--|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Foods or beverages are not offered at school celebrations | 5.8 | 3.1 - 10.7 | 133 | 0.5 | 0.1 - 2.3 | 206 | 0.0 | † | 19 | 2.5 | 1.4 - 4.4 | 358 |
| b. Never | 0.7 | 0.1 - 3.5 | 133 | 2.0 | 0.9 - 4.4 | 206 | 0.0 | † | 19 | 1.3 | 0.7 - 2.7 | 358 |
| c. Rarely | 13.3 | 9.2 - 18.8 | 133 | 10.5 | 7.5 - 14.4 | 206 | 0.0 | † | 19 | 10.8 | 8.4 - 13.7 | 358 |
| d. Sometimes | 58.1 | 50.6 - 65.3 | 133 | 57.8 | 52.9 - 62.6 | 206 | 69.4 | 46.4 - 85.6 | 19 | 58.8 | 54.6 - 62.9 | 358 |
| e. Always or almost always | 22.1 | 16.7 - 28.5 | 133 | 29.3 | 24.8 - 34.1 | 206 | 30.6 | 14.4 - 53.6 | 19 | 26.6 | 23.1 - 30.5 | 358 |

† Confidence intervals not applicable for 0 percent or 100 percent.
N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

34. Percentage of schools in which students can purchase snack foods or beverages from one or more vending machines at the school or at a school store, canteen, or snack bar.

| High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---------------------|-------------------------------|-----|-----------------------|-------------------------------|-----|-----------------------------------|-------------------------------|----|--------------------|-------------------------------|-----|
| Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| 88.0 | 82.0 - 92.2 | 132 | 50.5 | 45.2 - 55.7 | 205 | 55.4 | 34.0 - 75.0 | 19 | 65.0 | 61.1 - 68.8 | 356 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

35. Percentage of schools in which students can purchase the following snack foods or beverages from vending machines or at the school store, canteen, or snack bar.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Chocolate candy | 42.8 | 35.4 - 50.5 | 131 | 15.0 | 11.6 - 19.0 | 205 | 40.1 | 24.6 - 57.9 | 19 | 27.3 | 23.7 - 31.1 | 355 |
| b. Other kinds of candy | 47.8 | 40.3 - 55.4 | 131 | 18.9 | 15.3 - 23.1 | 205 | 28.4 | 18.8 - 40.6 | 19 | 30.5 | 26.9 - 34.2 | 355 |
| c. Salty snacks that are not low in fat (e.g., regular potato chips) | 60.1 | 53.3 - 66.5 | 131 | 24.6 | 20.3 - 29.6 | 205 | 46.0 | 27.9 - 65.2 | 19 | 39.5 | 35.7 - 43.4 | 355 |
| d. Cookies, crackers, cakes, pastries, or other baked goods that are not low in fat | 57.6 | 50.3 - 64.6 | 131 | 23.7 | 19.1 - 29.0 | 205 | 51.8 | 31.0 - 72.0 | 19 | 38.5 | 34.4 - 42.7 | 355 |
| e. Ice cream or frozen yogurt that is not low in fat | 19.2 | 14.0 - 25.7 | 131 | 16.1 | 12.3 - 20.8 | 204 | 15.3 | 5.6 - 35.6 | 19 | 17.2 | 14.1 - 20.8 | 354 |
| f. 2% or whole milk (plain or flavored) | 42.1 | 36.5 - 47.9 | 128 | 24.9 | 20.3 - 30.1 | 204 | 19.0 | 7.8 - 39.2 | 19 | 30.8 | 27.3 - 34.6 | 351 |
| g. Water ices or frozen slushes that do not contain juice | 22.8 | 16.9 - 30.0 | 130 | 9.7 | 7.0 - 13.3 | 204 | 0.0 | † | 19 | 13.9 | 11.1 - 17.1 | 353 |
| h. Soda pop or fruit drinks that are not 100% juice | 54.7 | 47.5 - 61.8 | 131 | 18.0 | 14.6 - 22.0 | 205 | 37.9 | 21.9 - 57.0 | 18 | 33.2 | 29.7 - 37.0 | 354 |
| i. Sports drinks (e.g., Gatorade) | 78.7 | 71.7 - 84.3 | 130 | 33.4 | 28.5 - 38.7 | 205 | 46.0 | 27.9 - 65.2 | 19 | 51.3 | 47.3 - 55.2 | 354 |
| j. Foods or beverages containing caffeine | 49.5 | 42.9 - 56.2 | 131 | 16.9 | 13.5 - 20.9 | 205 | 15.3 | 5.6 - 35.6 | 19 | 29.0 | 25.8 - 32.5 | 355 |
| k. Fruits (not fruit juice) | 35.9 | 29.3 - 43.0 | 131 | 22.2 | 18.1 - 26.9 | 205 | 15.3 | 5.6 - 35.6 | 19 | 26.8 | 23.3 - 30.7 | 355 |
| l. Non-fried vegetables (not vegetable juice) | 32.7 | 26.4 - 39.7 | 131 | 17.4 | 13.7 - 21.7 | 205 | 9.5 | 2.8 - 27.5 | 19 | 22.5 | 19.3 - 26.2 | 355 |

† Confidence intervals not applicable for 0 percent or 100 percent.
N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

36. Percentage of schools that limit the package or serving size of any individual food and beverage items sold in vending machines or at the school store, canteen, or snack bar.

| High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---------------------|-------------------------|-----|-----------------------|-------------------------|-----|-----------------------------------|-------------------------|----|--------------------|-------------------------|-----|
| Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| 58.5 | 50.8 - 65.7 | 130 | 33.4 | 28.4 - 38.8 | 205 | 17.5 | 6.5 - 39.5 | 19 | 41.6 | 37.5 - 45.8 | 354 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

37. Percentage of schools that have done any of the following activities during this school year.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Priced nutritious foods and beverages at a lower cost while increasing the price of less nutritious foods and beverages | 9.7 | 6.1 - 15.0 | 132 | 5.4 | 3.3 - 8.6 | 205 | 0.0 | † | 19 | 6.6 | 4.7 - 9.1 | 356 |
| b. Collected suggestions from students, families, and school staff on nutritious food preferences and strategies to promote healthy eating | 37.2 | 30.1 - 44.8 | 132 | 30.8 | 25.9 - 36.1 | 206 | 27.0 | 15.1 - 43.5 | 19 | 32.9 | 28.9 - 37.1 | 357 |
| c. Provided information to students or families on the nutrition and caloric content of foods available | 42.6 | 36.0 - 49.4 | 132 | 44.1 | 39.0 - 49.4 | 206 | 15.3 | 10.4 - 22.1 | 19 | 41.4 | 37.6 - 45.3 | 357 |
| d. Conducted taste tests to determine food preferences for nutritious items | 18.9 | 14.0 - 25.1 | 132 | 15.6 | 11.8 - 20.4 | 205 | 9.5 | 2.8 - 27.5 | 19 | 16.4 | 13.4 - 19.9 | 356 |
| e. Provided opportunities for students to visit the cafeteria to learn about food safety, food preparation, or other nutrition-related topics | 15.4 | 10.9 - 21.3 | 132 | 11.3 | 8.0 - 15.7 | 205 | 13.1 | 5.4 - 28.4 | 19 | 13.0 | 10.3 - 16.2 | 356 |

† Confidence intervals not applicable for 0 percent or 100 percent.

N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

38. Percentage of schools that promote candy, meals from fast food restaurants, or soft drinks through the distribution of products, such as t-shirts, hats, and book covers to students.

| High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| 4.1 | 1.9 - 8.4 | 129 | 1.5 | 0.6 - 3.7 | 205 | 0.0 | † | 19 | 2.3 | 1.3 - 4.1 | 353 |

† Confidence intervals not applicable for 0 percent or 100 percent.
 N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

39. Percentage of schools that prohibit advertisements for candy, fast food restaurants, or soft drinks in the following locations.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. In the school building | 62.0 | 55.3 - 68.2 | 132 | 72.2 | 67.1 - 76.8 | 206 | 73.0 | 50.0 - 88.0 | 19 | 68.4 | 64.4 - 72.2 | 357 |
| b. On school grounds including on the outside of the school building, on playing fields, or other areas of the campus | 40.2 | 34.2 - 46.5 | 132 | 66.0 | 60.9 - 70.7 | 206 | 73.0 | 50.0 - 88.0 | 19 | 56.8 | 52.9 - 60.6 | 357 |
| c. On school buses or other vehicles used to transport students | 73.4 | 66.1 - 79.6 | 132 | 79.9 | 75.5 - 83.7 | 206 | 82.5 | 60.5 - 93.5 | 19 | 77.6 | 73.8 - 81.0 | 357 |
| d. In school publications (e.g., newsletters, newspapers, web sites, or other school publications) | 45.5 | 38.3 - 52.8 | 132 | 73.7 | 68.9 - 78.0 | 206 | 78.8 | 57.0 - 91.3 | 19 | 63.4 | 59.4 - 67.2 | 357 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

40. Percentage of schools that have a full-time registered nurse who provides health services to students.

| High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| 26.9 | 20.8 - 33.9 | 133 | 22.3 | 17.6 - 27.7 | 206 | 24.8 | 11.1 - 46.7 | 19 | 24.2 | 20.5 - 28.4 | 358 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

41. Percentage of schools that have an asthma action plan on file for all, most, some, or no students with known asthma.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|--|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. This school has no students with known asthma. | 0.0 | † | 131 | 0.0 | † | 205 | 5.8 | 1.1 - 25.4 | 19 | 0.4 | 0.1 - 2.1 | 355 |
| b. All students with known asthma have an asthma action plan on file. | 71.2 | 64.5 - 77.0 | 131 | 75.8 | 70.9 - 80.1 | 205 | 55.4 | 38.2 - 71.4 | 19 | 72.5 | 68.7 - 76.1 | 355 |
| c. Most students with known asthma have an asthma action plan on file. | 18.4 | 13.5 - 24.5 | 131 | 15.5 | 11.7 - 20.2 | 205 | 27.0 | 15.1 - 43.5 | 19 | 17.4 | 14.4 - 21.0 | 355 |
| d. Some students with known asthma have an asthma action plan on file. | 7.2 | 4.3 - 11.7 | 131 | 7.7 | 5.2 - 11.5 | 205 | 5.8 | 1.1 - 25.4 | 19 | 7.4 | 5.4 - 10.0 | 355 |
| e. No students with known asthma have an asthma action plan on file. | 3.3 | 1.4 - 7.4 | 131 | 1.0 | 0.3 - 3.0 | 205 | 5.8 | 1.1 - 25.4 | 19 | 2.2 | 1.2 - 4.1 | 355 |

† Confidence intervals not applicable for 0 percent or 100 percent.
N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

42. Percentage of schools that use the following events to identify students with poorly controlled asthma.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|--|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. This school does not identify students with poorly controlled asthma. | 15.4 | 10.9 - 21.3 | 132 | 13.1 | 9.8 - 17.3 | 200 | 5.8 | 1.1 - 25.4 | 19 | 13.4 | 10.8 - 16.6 | 351 |
| b. Frequent absences from school | 42.7 | 35.2 - 50.6 | 132 | 49.1 | 43.8 - 54.4 | 201 | 34.3 | 17.4 - 56.4 | 19 | 45.5 | 41.2 - 50.0 | 352 |
| c. Frequent visits to the school health office due to asthma | 62.5 | 56.2 - 68.4 | 132 | 68.4 | 63.5 - 72.9 | 203 | 49.6 | 31.0 - 68.3 | 19 | 64.8 | 60.9 - 68.4 | 354 |
| d. Frequent asthma symptoms at school | 58.5 | 51.7 - 65.0 | 133 | 68.6 | 63.3 - 73.4 | 202 | 59.1 | 42.4 - 73.9 | 19 | 64.0 | 60.0 - 67.9 | 354 |
| e. Frequent non-participation in physical education class due to asthma | 44.1 | 37.1 - 51.4 | 133 | 49.6 | 44.6 - 54.7 | 201 | 53.2 | 34.9 - 70.7 | 19 | 47.8 | 43.7 - 51.9 | 353 |
| f. Students sent home early due to asthma | 47.2 | 39.9 - 54.6 | 133 | 53.4 | 47.7 - 59.0 | 200 | 53.2 | 53.2 - 53.2 | 19 | 51.0 | 46.8 - 55.2 | 352 |
| g. Calls from school to 911, or other local emergency numbers, due to asthma | 42.3 | 34.8 - 50.1 | 133 | 35.0 | 29.9 - 40.5 | 202 | 43.7 | 27.8 - 61.0 | 19 | 38.4 | 34.2 - 42.8 | 354 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

43. Percentage of schools that provide each of the following services for students with poorly controlled asthma.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|--|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Providing referrals to primary healthcare clinicians or child health insurance programs | 61.4 | 54.3 - 68.0 | 132 | 67.7 | 62.7 - 72.4 | 202 | - | - | 18 | 65.1 | 61.2 - 68.9 | 352 |
| b. Ensuring an appropriate written asthma action plan is obtained | 82.7 | 77.2 - 87.1 | 133 | 91.2 | 87.2 - 94.0 | 206 | - | - | 18 | 87.7 | 84.8 - 90.1 | 357 |
| c. Ensuring access to and appropriate use of asthma medications, spacers, and peak flow meters at school | 79.5 | 73.5 - 84.4 | 133 | 90.5 | 86.4 - 93.5 | 203 | - | - | 18 | 85.7 | 82.5 - 88.3 | 354 |
| d. Offering asthma education for students with asthma | 54.1 | 46.8 - 61.2 | 132 | 60.7 | 55.3 - 65.8 | 204 | - | - | 18 | 57.3 | 53.2 - 61.4 | 354 |
| e. Minimizing asthma triggers in the school environment | 70.3 | 63.5 - 76.3 | 133 | 77.8 | 72.8 - 82.1 | 204 | - | - | 17 | 73.3 | 69.4 - 76.9 | 354 |
| f. Addressing social and emotional issues related to asthma | 50.2 | 42.9 - 57.5 | 133 | 58.3 | 52.9 - 63.5 | 203 | - | - | 17 | 53.9 | 49.6 - 58.1 | 353 |
| g. Providing additional psychosocial counseling or support services as needed | 56.0 | 49.1 - 62.7 | 133 | 59.2 | 54.0 - 64.2 | 202 | - | - | 17 | 56.6 | 52.6 - 60.6 | 352 |
| h. Ensuring access to safe, enjoyable physical education and activity opportunities | 87.4 | 82.0 - 91.4 | 133 | 92.6 | 89.1 - 95.0 | 205 | 70.8 | 60.6 - 79.2 | 19 | 89.0 | 86.3 - 91.2 | 357 |
| i. Ensuring access to preventive medications before physical activity | 84.5 | 78.5 - 89.0 | 132 | 88.5 | 84.6 - 91.6 | 204 | - | - | 18 | 86.0 | 83.1 - 88.6 | 354 |

- Results are suppressed due to insufficient number of respondents in subgroup.

N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

44. Percentage of schools in which school staff members are required to receive training on recognizing and responding to severe asthma symptoms more than once per year, once per year, or less than once per year.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|----------------------------|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. More than once per year | 2.8 | 1.3 - 6.2 | 133 | 1.9 | 0.9 - 4.3 | 206 | 3.6 | 0.7 - 16.6 | 19 | 2.4 | 1.4 - 4.1 | 358 |
| b. Once per year | 47.8 | 40.4 - 55.3 | 133 | 58.3 | 52.9 - 63.6 | 206 | 46.0 | 27.9 - 65.2 | 19 | 53.4 | 49.1 - 57.8 | 358 |
| c. Less than once per year | 4.0 | 1.9 - 8.1 | 133 | 8.6 | 6.0 - 12.2 | 206 | 0.0 | † | 19 | 6.2 | 4.5 - 8.6 | 358 |
| d. No such requirement | 45.4 | 38.6 - 52.4 | 133 | 31.1 | 26.3 - 36.4 | 206 | 50.4 | 31.7 - 69.0 | 19 | 38.0 | 34.0 - 42.1 | 358 |

† Confidence intervals not applicable for 0 percent or 100 percent.
N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

45. Percentage of schools that have adopted a policy stating that students are permitted to carry and self-administer asthma medications.

| High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---------------------|-------------------------------|-----|-----------------------|-------------------------------|-----|-----------------------------------|-------------------------------|----|--------------------|-------------------------------|-----|
| Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| 84.3 | 78.6 - 88.7 | 132 | 84.3 | 79.8 - 87.9 | 206 | 82.5 | 60.5 - 93.5 | 19 | 84.2 | 80.8 - 87.1 | 357 |

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

46. Percentage of schools that have procedures to inform each of the following groups about the policy permitting students to carry and self-administer asthma medications.*

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|-------------------------|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. Students | 95.7 | 91.4 - 97.9 | 112 | 94.2 | 90.5 - 96.5 | 171 | 100.0 | † | 16 | 95.2 | 92.8 - 96.8 | 299 |
| b. Parents and families | 94.9 | 90.3 - 97.3 | 112 | 92.3 | 88.1 - 95.1 | 169 | 100.0 | † | 16 | 93.8 | 91.2 - 95.7 | 297 |

*Among schools that have adopted a policy stating that students are permitted to carry and self-administer asthma medications.

† Confidence intervals not applicable for 0 percent or 100 percent.

N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

47. Percentage of schools that have the following person responsible for implementing the policy permitting students to carry and self-administer asthma medication.*

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. No single individual is responsible | 20.0 | 14.8 - 26.5 | 103 | 17.8 | 13.5 - 23.1 | 163 | 24.7 | 11.2 - 46.2 | 15 | 19.1 | 15.7 - 23.1 | 281 |
| b. Principal | 8.2 | 5.1 - 13.0 | 103 | 12.7 | 9.0 - 17.8 | 163 | 9.5 | 2.8 - 27.4 | 15 | 10.8 | 8.2 - 14.2 | 281 |
| c. Assistant principal | 0.9 | 0.2 - 4.5 | 103 | 0.6 | 0.1 - 3.2 | 163 | 0.0 | † | 15 | 0.7 | 0.2 - 2.2 | 281 |
| d. School nurse | 61.0 | 53.3 - 68.3 | 103 | 60.0 | 53.0 - 66.6 | 163 | 50.5 | 29.3 - 71.5 | 15 | 59.7 | 54.6 - 64.6 | 281 |
| e. Other school faculty or staff member | 9.8 | 5.9 - 16.0 | 103 | 8.8 | 5.7 - 13.3 | 163 | 15.3 | 4.8 - 38.9 | 15 | 9.7 | 7.0 - 13.1 | 281 |

*Among schools that have adopted a policy stating that students are permitted to carry and self-administer asthma medications.

† Confidence intervals not applicable for 0 percent or 100 percent.

N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

48. Percentage of schools in which students' families helped develop or implement policies and programs related to the following topics during the past two years.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. HIV, STD, or teen pregnancy prevention | 17.7 | 13.3 - 23.1 | 132 | 10.8 | 7.6 - 15.0 | 204 | 0.0 | † | 19 | 12.6 | 10.1 - 15.6 | 355 |
| b. Tobacco-use prevention | 30.4 | 24.5 - 37.0 | 133 | 25.0 | 20.4 - 30.1 | 205 | 0.0 | † | 19 | 25.2 | 21.8 - 28.9 | 357 |
| c. Physical activity | 20.5 | 15.0 - 27.4 | 132 | 28.0 | 23.0 - 33.6 | 205 | 5.8 | 1.1 - 25.4 | 19 | 23.5 | 20.0 - 27.5 | 356 |
| d. Nutrition and healthy eating | 28.1 | 22.2 - 34.7 | 132 | 28.9 | 23.9 - 34.5 | 205 | 15.3 | 5.6 - 35.6 | 19 | 27.6 | 23.9 - 31.6 | 356 |
| e. Asthma | 16.2 | 12.1 - 21.4 | 132 | 17.8 | 14.1 - 22.1 | 204 | 5.8 | 1.1 - 25.4 | 19 | 16.3 | 13.6 - 19.4 | 355 |

† Confidence intervals not applicable for 0 percent or 100 percent.
N = Unweighted number of observations

NORTH CAROLINA
2010 School Health Profiles Report
Weighted Principal Survey Results

49. Percentage of schools in which community members helped develop or implement policies and programs related to the following topics during the past two years.

| | High Schools | | | Middle Schools | | | Junior/Senior High Schools | | | All Schools | | |
|---|--------------|-------------------------|-----|----------------|-------------------------|-----|----------------------------|-------------------------|----|-------------|-------------------------|-----|
| | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N | Percent | 95% Confidence Interval | N |
| a. HIV, STD, or teen pregnancy prevention | 24.6 | 19.2 - 30.9 | 133 | 25.8 | 20.9 - 31.3 | 205 | 44.6 | 28.6 - 61.8 | 19 | 26.7 | 23.0 - 30.7 | 357 |
| b. Tobacco-use prevention | 40.3 | 33.8 - 47.1 | 133 | 37.1 | 31.9 - 42.7 | 205 | 38.7 | 21.6 - 59.1 | 19 | 38.4 | 34.4 - 42.7 | 357 |
| c. Physical activity | 23.9 | 18.2 - 30.7 | 133 | 35.7 | 30.4 - 41.4 | 205 | 42.3 | 24.7 - 62.1 | 19 | 31.7 | 27.8 - 36.0 | 357 |
| d. Nutrition and healthy eating | 34.2 | 28.0 - 41.0 | 133 | 41.8 | 36.3 - 47.5 | 206 | 44.6 | 28.6 - 61.8 | 19 | 39.1 | 35.1 - 43.3 | 358 |
| e. Asthma | 21.9 | 17.2 - 27.6 | 132 | 23.4 | 19.3 - 28.1 | 205 | 29.2 | 15.9 - 47.5 | 19 | 23.3 | 20.1 - 26.8 | 356 |