

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Total												
Injury and Violence												
Among students who rode a bicycle during the past 12 months, the percentage who never or rarely wore a bicycle helmet							83.2	84.7	81.7	No	No	
Percentage of students who never or rarely wore a seat belt when riding in a car driven by someone else							9.8	7.4	9.2	No	No	
Percentage of students who rode one or more times during the past 30 days in a car or other vehicle driven by someone who had been drinking alcohol							18.8	23.0	20.5	No	No	
Percentage of students who drove a car or other vehicle one or more times during the past 30 days when they had been drinking alcohol							7.2	10.2	10.0	No	No	
Percentage of students who carried a weapon such as a gun, knife, or club on one or more of the past 30 days							25.8	25.7	24.1	No	No	
Percentage of students who carried a weapon such as a gun, knife, or club on school property on one or more of the past 30 days							8.8	8.7	7.3	No	No	
Percentage of students who did not go to school on one or more of the past 30 days because they felt they would be unsafe at school or on their way to or from school							2.4	3.7	5.5	No	No	

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Total												
Injury and Violence												
Percentage of students who had been threatened or injured with a weapon such as a gun, knife, or club on school property one or more times during the past 12 months							7.2	6.8	8.4	No	No	
Percentage of students who had property, such as their car, clothing, or books stolen or deliberately damaged on school property one or more times during the past 12 months							26.8	26.2	29.5	No	No	
Percentage of students who were in a physical fight one or more times during the past 12 months							28.8	27.8	30.5	No	No	
Percentage of students who were in a physical fight one or more times during the past 12 months in which they were injured and had to be treated by a doctor or nurse							3.5	3.5	4.4	No	No	
Percentage of students who were in a physical fight on school property one or more times during the past 12 months							12.9	11.9	11.5	No	No	
Percentage of students who were ever hit, slapped, or physically hurt on purpose by their boyfriend or girlfriend during the past 12 months								11.6	13.9	No		
Percentage of students who had ever been physically forced to have sexual intercourse when they did not want to								9.4	8.4	No		

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Total												
Injury and Violence												
Percentage of students who felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months							29.1	28.3	28.9		No	No
Percentage of students who seriously considered attempting suicide during the past 12 months							19.3	16.1	13.0		Yes	No
Percentage of students who made a plan about how they would attempt suicide during the past 12 months								12.8	11.3		No	
Percentage of students who actually attempted suicide one or more times during the past 12 months								14.0	12.5		No	

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Total												
Tobacco Use												
Percentage of students who smoked a whole cigarette for the first time before age 13 years							23.4	20.0	19.5		No	No
Percentage of students who smoked cigarettes on one or more of the past 30 days							26.0	25.8	23.9		No	No
Percentage of students who smoked cigarettes on 20 or more of the past 30 days							11.8	12.3	8.2		Yes	No
Among students who reported current cigarette use, the percentage who ever tried to quit smoking cigarettes during the past 12 months							52.5	52.7	51.2		No	No

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Total												
Alcohol and Other Drug Use												
Percentage of students who had their first drink of alcohol other than a few sips before age 13 years							20.5	21.5	20.8		No	No
Percentage of students who had at least one drink of alcohol on one or more of the past 30 days							38.7	40.9	39.2		No	No
Percentage of students who had five or more drinks of alcohol in a row, that is, within a couple of hours, on one or more of the past 30 days							22.5	23.9	25.3		No	No
Percentage of students who had at least one drink of alcohol on school property on one or more of the past 30 days							4.4	5.1	5.6		No	No
Percentage of students who used marijuana one or more times during their life							37.2	36.4	35.0		No	No
Percentage of students who tried marijuana for the first time before age 13 years							8.6	9.2	9.3		No	No
Percentage of students who used marijuana one or more times during the past 30 days							18.5	18.8	17.0		No	No
Percentage of students who used marijuana on school property one or more times during the past 30 days							3.9	4.2	4.5		No	No
Percentage of students who used any form of cocaine, including powder, crack, or freebase one or more times during their life							7.5	9.0	10.7		No	No

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Total												
Alcohol and Other Drug Use												
Percentage of students who sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high one or more times during their life							16.2	15.7	15.9	No	No	
Percentage of students who used heroin one or more times during their life							3.2	3.3	4.9	No	No	
Percentage of students who used methamphetamines one or more times during their life							8.9	8.0	8.0	No	No	
Percentage of students who used ecstasy one or more times during their life							6.9	8.5	9.0	No	No	
Percentage of students who took steroid pills or shots without a doctor's prescription one or more times during their life							6.4	4.8	6.3	No	No	
Percentage of students who used a needle to inject any illegal drug into their body one or more times during their life							2.7	2.9	2.3	No	No	
Percentage of students who were offered, sold, or given an illegal drug by someone on school property during the past 12 months							31.6	25.8	27.9	No	Yes	

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Total												
Sexual Behaviors												
Percentage of students who ever had sexual intercourse							44.9	48.1	47.4		No	No
Percentage of students who had sexual intercourse for the first time before age 13 years							6.0	6.7	7.0		No	No
Percentage of students who had sexual intercourse with four or more people during their life							12.1	13.9	12.2		No	No
Percentage of students who had sexual intercourse with one or more people during the past three months							34.3	34.9	33.5		No	No
Among students who had sexual intercourse during the past three months, the percentage who drank alcohol or used drugs before last sexual intercourse							17.2	21.0	22.2		No	No
Among students who had sexual intercourse during the past three months, the percentage who used a condom during last sexual intercourse							53.0	59.1	57.8		No	No
Among students who had sexual intercourse during the past three months, the percentage who used birth control pills to prevent pregnancy before last sexual intercourse							19.2	22.8	23.2		No	No

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Total												
Weight Management and Dietary Behaviors												
Percentage of students who were at risk for becoming overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)**							13.6	15.3	15.4		No	No
Percentage of students who were overweight (i.e., at or above the 95th percentile for body mass index, by age and sex)**							15.2	14.3	12.5		No	No
Percentage of students who described themselves as slightly or very overweight							33.5	32.8	26.6		Yes	No
Percentage of students who were trying to lose weight							47.3	47.8	46.8		No	No
Percentage of students who exercised to lose weight or to keep from gaining weight during the past 30 days							63.5	64.1	65.9		No	No
Percentage of students who ate less food, fewer calories, or foods low in fat to lose weight or to keep from gaining weight during the past 30 days								42.0	43.2		No	
Percentage of students who took any diet pills, powders, or liquids without a doctor's advice to lose weight or to keep from gaining weight during the past 30 days							11.1	7.4	9.0		No	Yes

**Overweight and at risk for becoming overweight prevalence estimates for 1999 differ slightly from previously published results because different BMI cut points were used in 1999 than in subsequent years. To make these prevalence estimates comparable, the 1999 prevalence estimates were recalculated using the updated BMI cut points.

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Total												
Weight Management and Dietary Behaviors												
Percentage of students who vomited or took laxatives to lose weight or to keep from gaining weight during the past 30 days							3.8	5.1	6.2	Yes	No	
Percentage of students who drank three or more glasses per day of milk during the past seven days							11.9	14.7	10.0	No	Yes	

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year										Linear Change	Quadratic Change
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Total												
Physical Activity												
Percentage of students who were physically active for a total of at least 60 minutes per day on five or more of the past seven days								50.1	46.1		No	
Percentage of students who watched three or more hours per day of TV on an average school day								30.5	26.4		Yes	
Percentage of students who attended physical education (PE) classes on one or more days in an average week when they were in school							49.1	40.4	45.1		No	No
Percentage of students who attended physical education (PE) classes daily in an average week when they were in school							38.3	32.1	35.7		No	No

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year										Linear Change	Quadratic Change
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Total												
Other												
Percentage of students who had ever been told by a doctor or nurse that they had asthma								19.6	20.6		No	

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Male												
Injury and Violence												
Among students who rode a bicycle during the past 12 months, the percentage who never or rarely wore a bicycle helmet							84.4	85.9	83.1	No	No	
Percentage of students who never or rarely wore a seat belt when riding in a car driven by someone else							12.6	8.8	13.6	No	Yes	
Percentage of students who rode one or more times during the past 30 days in a car or other vehicle driven by someone who had been drinking alcohol							19.5	22.8	23.1	No	No	
Percentage of students who drove a car or other vehicle one or more times during the past 30 days when they had been drinking alcohol							9.9	12.5	12.5	No	No	
Percentage of students who carried a weapon such as a gun, knife, or club on one or more of the past 30 days							42.2	41.4	38.6	No	No	
Percentage of students who carried a weapon such as a gun, knife, or club on school property on one or more of the past 30 days							14.7	13.8	11.8	No	No	
Percentage of students who did not go to school on one or more of the past 30 days because they felt they would be unsafe at school or on their way to or from school							3.3	3.4	5.9	No	No	

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Male												
Injury and Violence												
Percentage of students who had been threatened or injured with a weapon such as a gun, knife, or club on school property one or more times during the past 12 months							8.9	9.1	9.7	No	No	
Percentage of students who had property, such as their car, clothing, or books stolen or deliberately damaged on school property one or more times during the past 12 months							29.5	30.7	31.5	No	No	
Percentage of students who were in a physical fight one or more times during the past 12 months							37.4	35.1	40.6	No	No	
Percentage of students who were in a physical fight one or more times during the past 12 months in which they were injured and had to be treated by a doctor or nurse							5.2	4.9	4.6	No	No	
Percentage of students who were in a physical fight on school property one or more times during the past 12 months							17.4	15.2	16.5	No	No	
Percentage of students who were ever hit, slapped, or physically hurt on purpose by their boyfriend or girlfriend during the past 12 months								10.4	15.9	Yes		
Percentage of students who had ever been physically forced to have sexual intercourse when they did not want to								6.2	6.2	No		

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Male												
Injury and Violence												
Percentage of students who felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months							24.0	19.3	23.6	No	No	
Percentage of students who seriously considered attempting suicide during the past 12 months							17.8	10.4	11.4	Yes	Yes	
Percentage of students who made a plan about how they would attempt suicide during the past 12 months								8.4	11.1	No		
Percentage of students who actually attempted suicide one or more times during the past 12 months								14.6	14.5	No		

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Male												
Tobacco Use												
Percentage of students who smoked a whole cigarette for the first time before age 13 years							28.3	21.4	22.4	No	No	
Percentage of students who smoked cigarettes on one or more of the past 30 days							27.2	26.1	22.7	No	No	
Percentage of students who smoked cigarettes on 20 or more of the past 30 days							12.7	12.1	7.0	Yes	No	
Among students who reported current cigarette use, the percentage who ever tried to quit smoking cigarettes during the past 12 months								42.8	47.4	No		

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Male												
Alcohol and Other Drug Use												
Percentage of students who had their first drink of alcohol other than a few sips before age 13 years							26.1	26.4	25.3	No	No	
Percentage of students who had at least one drink of alcohol on one or more of the past 30 days							43.3	42.9	39.9	No	No	
Percentage of students who had five or more drinks of alcohol in a row, that is, within a couple of hours, on one or more of the past 30 days							28.6	28.0	25.5	No	No	
Percentage of students who had at least one drink of alcohol on school property on one or more of the past 30 days							6.4	7.4	8.5	No	No	
Percentage of students who used marijuana one or more times during their life							41.9	39.5	38.9	No	No	
Percentage of students who tried marijuana for the first time before age 13 years							13.6	11.3	10.8	No	No	
Percentage of students who used marijuana one or more times during the past 30 days							21.8	21.7	19.0	No	No	
Percentage of students who used marijuana on school property one or more times during the past 30 days							6.5	6.0	5.9	No	No	
Percentage of students who used any form of cocaine, including powder, crack, or freebase one or more times during their life							8.5	10.4	11.6	No	No	

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Male												
Alcohol and Other Drug Use												
Percentage of students who sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high one or more times during their life							16.5	17.2	15.6		No	No
Percentage of students who used heroin one or more times during their life							4.1	5.0	7.2		No	No
Percentage of students who used methamphetamines one or more times during their life							10.2	9.1	8.9		No	No
Percentage of students who used ecstasy one or more times during their life							8.9	10.2	10.3		No	No
Percentage of students who took steroid pills or shots without a doctor's prescription one or more times during their life							8.2	6.4	8.9		No	No
Percentage of students who used a needle to inject any illegal drug into their body one or more times during their life							3.7	3.7	3.8		No	No
Percentage of students who were offered, sold, or given an illegal drug by someone on school property during the past 12 months							35.0	28.5	33.5		No	Yes

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Male												
Sexual Behaviors												
Percentage of students who ever had sexual intercourse							46.1	46.7	48.1	No	No	
Percentage of students who had sexual intercourse for the first time before age 13 years							10.9	8.6	8.8	No	No	
Percentage of students who had sexual intercourse with four or more people during their life							15.4	15.6	13.2	No	No	
Percentage of students who had sexual intercourse with one or more people during the past three months							33.9	33.5	31.1	No	No	
Among students who had sexual intercourse during the past three months, the percentage who drank alcohol or used drugs before last sexual intercourse							24.3	26.7	27.8	No	No	
Among students who had sexual intercourse during the past three months, the percentage who used a condom during last sexual intercourse								60.6	66.0	No		
Among students who had sexual intercourse during the past three months, the percentage who used birth control pills to prevent pregnancy before last sexual intercourse								24.0	17.3	No		

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Male												
Weight Management and Dietary Behaviors												
Percentage of students who were at risk for becoming overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)**							14.7	15.1	17.1	No	No	
Percentage of students who were overweight (i.e., at or above the 95th percentile for body mass index, by age and sex)**							19.6	16.4	16.4	No	No	
Percentage of students who described themselves as slightly or very overweight							29.0	24.4	24.1	No	No	
Percentage of students who were trying to lose weight							34.3	32.6	35.7	No	No	
Percentage of students who exercised to lose weight or to keep from gaining weight during the past 30 days							55.5	54.8	62.8	Yes	No	
Percentage of students who ate less food, fewer calories, or foods low in fat to lose weight or to keep from gaining weight during the past 30 days								27.8	32.5	Yes		
Percentage of students who took any diet pills, powders, or liquids without a doctor's advice to lose weight or to keep from gaining weight during the past 30 days							10.0	6.6	9.7	No	Yes	

**Overweight and at risk for becoming overweight prevalence estimates for 1999 differ slightly from previously published results because different BMI cut points were used in 1999 than in subsequent years. To make these prevalence estimates comparable, the 1999 prevalence estimates were recalculated using the updated BMI cut points.

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Male												
Weight Management and Dietary Behaviors												
Percentage of students who vomited or took laxatives to lose weight or to keep from gaining weight during the past 30 days							3.0	3.0	6.1	No	No	
Percentage of students who drank three or more glasses per day of milk during the past seven days							18.1	18.8	11.3	Yes	Yes	

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year										Linear Change	Quadratic Change
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Male												
Physical Activity												
Percentage of students who were physically active for a total of at least 60 minutes per day on five or more of the past seven days								60.1	57.8		No	
Percentage of students who watched three or more hours per day of TV on an average school day								32.8	30.9		No	
Percentage of students who attended physical education (PE) classes on one or more days in an average week when they were in school							59.1	46.8	54.5		No	No
Percentage of students who attended physical education (PE) classes daily in an average week when they were in school							45.3	36.5	43.0		No	No

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year										Linear Change	Quadratic Change
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Male												
Other												
Percentage of students who had ever been told by a doctor or nurse that they had asthma								20.3	19.8		No	

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change
	1991	1993	1995	1997	1999	2001	2003	2005	2007		
Female											
Injury and Violence											
Among students who rode a bicycle during the past 12 months, the percentage who never or rarely wore a bicycle helmet							81.4	82.8	80.6	No	No
Percentage of students who never or rarely wore a seat belt when riding in a car driven by someone else							6.8	6.0	4.2	No	No
Percentage of students who rode one or more times during the past 30 days in a car or other vehicle driven by someone who had been drinking alcohol							18.0	23.1	17.6	No	Yes
Percentage of students who drove a car or other vehicle one or more times during the past 30 days when they had been drinking alcohol							4.4	7.8	7.0	No	No
Percentage of students who carried a weapon such as a gun, knife, or club on one or more of the past 30 days							9.2	9.5	9.0	No	No
Percentage of students who carried a weapon such as a gun, knife, or club on school property on one or more of the past 30 days							2.8	3.4	2.9	No	No
Percentage of students who did not go to school on one or more of the past 30 days because they felt they would be unsafe at school or on their way to or from school							1.5	3.7	4.4	No	No

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Female												
Injury and Violence												
Percentage of students who had been threatened or injured with a weapon such as a gun, knife, or club on school property one or more times during the past 12 months							5.5	4.3	6.5	No	No	
Percentage of students who had property, such as their car, clothing, or books stolen or deliberately damaged on school property one or more times during the past 12 months							23.9	21.6	27.1	No	No	
Percentage of students who were in a physical fight one or more times during the past 12 months							20.0	20.4	20.0	No	No	
Percentage of students who were in a physical fight one or more times during the past 12 months in which they were injured and had to be treated by a doctor or nurse							1.7	2.0	3.6	No	No	
Percentage of students who were in a physical fight on school property one or more times during the past 12 months							8.2	8.5	6.4	No	No	
Percentage of students who were ever hit, slapped, or physically hurt on purpose by their boyfriend or girlfriend during the past 12 months								13.0	11.9	No		
Percentage of students who had ever been physically forced to have sexual intercourse when they did not want to								12.8	10.6	No		

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change
	1991	1993	1995	1997	1999	2001	2003	2005	2007		
Female											
Injury and Violence											
Percentage of students who felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months							34.5	37.6	34.4	No	No
Percentage of students who seriously considered attempting suicide during the past 12 months							20.8	22.1	14.6	No	No
Percentage of students who made a plan about how they would attempt suicide during the past 12 months								17.3	11.5	Yes	
Percentage of students who actually attempted suicide one or more times during the past 12 months								13.2	10.5	No	

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year										Linear Change	Quadratic Change
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Female												
Tobacco Use												
Percentage of students who smoked a whole cigarette for the first time before age 13 years							18.5	18.4	16.0		No	No
Percentage of students who smoked cigarettes on one or more of the past 30 days							24.8	25.4	24.4		No	No
Percentage of students who smoked cigarettes on 20 or more of the past 30 days							10.9	12.6	8.9		No	No
Among students who reported current cigarette use, the percentage who ever tried to quit smoking cigarettes during the past 12 months							57.1	63.0	53.4		No	No

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Female												
Alcohol and Other Drug Use												
Percentage of students who had their first drink of alcohol other than a few sips before age 13 years							14.8	16.2	16.5		No	No
Percentage of students who had at least one drink of alcohol on one or more of the past 30 days							33.9	38.6	38.3		No	No
Percentage of students who had five or more drinks of alcohol in a row, that is, within a couple of hours, on one or more of the past 30 days							16.4	19.8	24.6		Yes	No
Percentage of students who had at least one drink of alcohol on school property on one or more of the past 30 days							2.4	2.3	2.3		No	No
Percentage of students who used marijuana one or more times during their life							32.6	33.3	30.6		No	No
Percentage of students who tried marijuana for the first time before age 13 years							3.5	6.8	7.1		No	No
Percentage of students who used marijuana one or more times during the past 30 days							15.2	15.9	14.7		No	No
Percentage of students who used marijuana on school property one or more times during the past 30 days							1.2	2.2	2.6		No	No
Percentage of students who used any form of cocaine, including powder, crack, or freebase one or more times during their life							6.5	7.4	9.3		No	No

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Female												
Alcohol and Other Drug Use												
Percentage of students who sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high one or more times during their life							15.9	14.2	15.6	No	No	
Percentage of students who used heroin one or more times during their life							2.3	1.5	2.0	No	No	
Percentage of students who used methamphetamines one or more times during their life							7.5	6.9	6.5	No	No	
Percentage of students who used ecstasy one or more times during their life							4.9	6.6	6.9	No	No	
Percentage of students who took steroid pills or shots without a doctor's prescription one or more times during their life							4.5	3.2	3.1	No	No	
Percentage of students who used a needle to inject any illegal drug into their body one or more times during their life							1.6	2.0	0.8	No	No	
Percentage of students who were offered, sold, or given an illegal drug by someone on school property during the past 12 months							28.0	23.0	22.6	No	No	

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Female												
Sexual Behaviors												
Percentage of students who ever had sexual intercourse							43.6	49.6	46.5	No	No	
Percentage of students who had sexual intercourse for the first time before age 13 years							1.1	4.8	5.3	Yes	No	
Percentage of students who had sexual intercourse with four or more people during their life							8.8	12.3	11.3	No	No	
Percentage of students who had sexual intercourse with one or more people during the past three months							34.8	36.4	35.4	No	No	
Among students who had sexual intercourse during the past three months, the percentage who drank alcohol or used drugs before last sexual intercourse							10.1	15.7	18.0	No	No	
Among students who had sexual intercourse during the past three months, the percentage who used a condom during last sexual intercourse							49.9	57.7	52.4	No	No	
Among students who had sexual intercourse during the past three months, the percentage who used birth control pills to prevent pregnancy before last sexual intercourse							23.5	21.8	28.5	No	No	

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Female												
Weight Management and Dietary Behaviors												
Percentage of students who were at risk for becoming overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)**							12.4	15.6	13.6		No	No
Percentage of students who were overweight (i.e., at or above the 95th percentile for body mass index, by age and sex)**							10.7	12.1	8.2		No	No
Percentage of students who described themselves as slightly or very overweight							38.2	41.4	29.5		Yes	Yes
Percentage of students who were trying to lose weight							60.7	63.7	58.3		No	No
Percentage of students who exercised to lose weight or to keep from gaining weight during the past 30 days							71.7	73.7	69.0		No	No
Percentage of students who ate less food, fewer calories, or foods low in fat to lose weight or to keep from gaining weight during the past 30 days								56.7	53.9		No	
Percentage of students who took any diet pills, powders, or liquids without a doctor's advice to lose weight or to keep from gaining weight during the past 30 days							12.3	8.2	8.2		No	No

**Overweight and at risk for becoming overweight prevalence estimates for 1999 differ slightly from previously published results because different BMI cut points were used in 1999 than in subsequent years. To make these prevalence estimates comparable, the 1999 prevalence estimates were recalculated using the updated BMI cut points.

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change
	1991	1993	1995	1997	1999	2001	2003	2005	2007		
Female											
Weight Management and Dietary Behaviors											
Percentage of students who vomited or took laxatives to lose weight or to keep from gaining weight during the past 30 days							4.6	7.3	6.3	No	No
Percentage of students who drank three or more glasses per day of milk during the past seven days							5.5	10.5	8.0	No	Yes

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year										Linear Change	Quadratic Change
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Female												
Physical Activity												
Percentage of students who were physically active for a total of at least 60 minutes per day on five or more of the past seven days								39.9	34.5		No	
Percentage of students who watched three or more hours per day of TV on an average school day								28.2	22.0		Yes	
Percentage of students who attended physical education (PE) classes on one or more days in an average week when they were in school								38.7	34.2	35.9	No	No
Percentage of students who attended physical education (PE) classes daily in an average week when they were in school								30.9	27.9	28.3	No	No

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year										Linear Change	Quadratic Change
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
Female												
Other												
Percentage of students who had ever been told by a doctor or nurse that they had asthma								18.7	21.3		No	

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
White*												
Injury and Violence												
Among students who rode a bicycle during the past 12 months, the percentage who never or rarely wore a bicycle helmet							81.9	83.6	81.6	No	No	
Percentage of students who never or rarely wore a seat belt when riding in a car driven by someone else							9.4	6.3	8.4	No	No	
Percentage of students who rode one or more times during the past 30 days in a car or other vehicle driven by someone who had been drinking alcohol							18.6	22.0	19.7	No	No	
Percentage of students who drove a car or other vehicle one or more times during the past 30 days when they had been drinking alcohol							7.8	9.6	9.8	No	No	
Percentage of students who carried a weapon such as a gun, knife, or club on one or more of the past 30 days							25.1	27.0	23.9	No	No	
Percentage of students who carried a weapon such as a gun, knife, or club on school property on one or more of the past 30 days							8.4	9.0	6.8	No	No	
Percentage of students who did not go to school on one or more of the past 30 days because they felt they would be unsafe at school or on their way to or from school							1.7	2.9	4.5	Yes	No	

*Non-Hispanic.

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
White*												
Injury and Violence												
Percentage of students who had been threatened or injured with a weapon such as a gun, knife, or club on school property one or more times during the past 12 months							6.6	6.4	7.3	No	No	
Percentage of students who had property, such as their car, clothing, or books stolen or deliberately damaged on school property one or more times during the past 12 months							26.8	26.0	29.4	No	No	
Percentage of students who were in a physical fight one or more times during the past 12 months							27.2	26.8	28.4	No	No	
Percentage of students who were in a physical fight one or more times during the past 12 months in which they were injured and had to be treated by a doctor or nurse							3.5	3.6	3.9	No	No	
Percentage of students who were in a physical fight on school property one or more times during the past 12 months							11.7	11.8	11.0	No	No	
Percentage of students who were ever hit, slapped, or physically hurt on purpose by their boyfriend or girlfriend during the past 12 months								11.3	13.4	No		
Percentage of students who had ever been physically forced to have sexual intercourse when they did not want to								8.1	8.5	No		

*Non-Hispanic.

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change
	1991	1993	1995	1997	1999	2001	2003	2005	2007		
White*											
Injury and Violence											
Percentage of students who felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months							29.6	26.7	28.5	No	No
Percentage of students who seriously considered attempting suicide during the past 12 months							19.5	14.6	12.7	Yes	No
Percentage of students who made a plan about how they would attempt suicide during the past 12 months								11.4	11.3	No	
Percentage of students who actually attempted suicide one or more times during the past 12 months								12.3	11.4	No	

*Non-Hispanic.

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change
	1991	1993	1995	1997	1999	2001	2003	2005	2007		
White*											
Tobacco Use											
Percentage of students who smoked a whole cigarette for the first time before age 13 years							24.5	19.3	19.1	Yes	No
Percentage of students who smoked cigarettes on one or more of the past 30 days							26.6	25.5	25.1	No	No
Percentage of students who smoked cigarettes on 20 or more of the past 30 days							12.2	12.6	8.6	No	No
Among students who reported current cigarette use, the percentage who ever tried to quit smoking cigarettes during the past 12 months							51.7	51.7	48.9	No	No

*Non-Hispanic.

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
White*												
Alcohol and Other Drug Use												
Percentage of students who had their first drink of alcohol other than a few sips before age 13 years							19.6	20.0	19.1	No	No	
Percentage of students who had at least one drink of alcohol on one or more of the past 30 days							38.6	40.1	40.9	No	No	
Percentage of students who had five or more drinks of alcohol in a row, that is, within a couple of hours, on one or more of the past 30 days							23.0	24.1	26.5	No	No	
Percentage of students who had at least one drink of alcohol on school property on one or more of the past 30 days							4.3	4.6	5.6	No	No	
Percentage of students who used marijuana one or more times during their life							36.1	34.9	33.7	No	No	
Percentage of students who tried marijuana for the first time before age 13 years							7.9	8.1	8.1	No	No	
Percentage of students who used marijuana one or more times during the past 30 days							17.8	17.3	17.0	No	No	
Percentage of students who used marijuana on school property one or more times during the past 30 days							3.6	3.7	4.4	No	No	
Percentage of students who used any form of cocaine, including powder, crack, or freebase one or more times during their life							8.0	9.2	9.8	No	No	

*Non-Hispanic.

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
White*												
Alcohol and Other Drug Use												
Percentage of students who sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high one or more times during their life							15.8	15.7	16.0	No	No	
Percentage of students who used heroin one or more times during their life							3.3	2.7	4.5	No	No	
Percentage of students who used methamphetamines one or more times during their life							9.5	7.9	7.6	No	No	
Percentage of students who used ecstasy one or more times during their life							6.9	7.9	8.5	No	No	
Percentage of students who took steroid pills or shots without a doctor's prescription one or more times during their life							6.3	4.1	5.7	No	No	
Percentage of students who used a needle to inject any illegal drug into their body one or more times during their life							2.9	2.5	1.9	No	No	
Percentage of students who were offered, sold, or given an illegal drug by someone on school property during the past 12 months							32.5	24.9	28.1	No	Yes	

*Non-Hispanic.

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
White*												
Sexual Behaviors												
Percentage of students who ever had sexual intercourse							42.9	44.5	45.1	No	No	
Percentage of students who had sexual intercourse for the first time before age 13 years							4.0	5.0	4.8	No	No	
Percentage of students who had sexual intercourse with four or more people during their life							9.8	10.6	10.4	No	No	
Percentage of students who had sexual intercourse with one or more people during the past three months							32.3	32.2	31.9	No	No	
Among students who had sexual intercourse during the past three months, the percentage who drank alcohol or used drugs before last sexual intercourse							19.6	19.5	22.7	No	No	
Among students who had sexual intercourse during the past three months, the percentage who used a condom during last sexual intercourse							51.8	59.4	56.5	No	No	
Among students who had sexual intercourse during the past three months, the percentage who used birth control pills to prevent pregnancy before last sexual intercourse							21.5	24.4	24.9	No	No	

*Non-Hispanic.

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change	
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
White*												
Weight Management and Dietary Behaviors												
Percentage of students who were at risk for becoming overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)**							13.4	14.5	13.9	No	No	
Percentage of students who were overweight (i.e., at or above the 95th percentile for body mass index, by age and sex)**							15.3	13.5	12.5	No	No	
Percentage of students who described themselves as slightly or very overweight							33.9	32.1	27.1	Yes	No	
Percentage of students who were trying to lose weight							47.5	47.6	47.6	No	No	
Percentage of students who exercised to lose weight or to keep from gaining weight during the past 30 days							64.9	65.0	66.6	No	No	
Percentage of students who ate less food, fewer calories, or foods low in fat to lose weight or to keep from gaining weight during the past 30 days								42.6	44.6	No		
Percentage of students who took any diet pills, powders, or liquids without a doctor's advice to lose weight or to keep from gaining weight during the past 30 days							11.3	7.5	9.1	No	Yes	

*Non-Hispanic.

**Overweight and at risk for becoming overweight prevalence estimates for 1999 differ slightly from previously published results because different BMI cut points were used in 1999 than in subsequent years. To make these prevalence estimates comparable, the 1999 prevalence estimates were recalculated using the updated BMI cut points.

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year									Linear Change	Quadratic Change
	1991	1993	1995	1997	1999	2001	2003	2005	2007		
White*											
Weight Management and Dietary Behaviors											
Percentage of students who vomited or took laxatives to lose weight or to keep from gaining weight during the past 30 days							3.7	4.1	6.1	No	No
Percentage of students who drank three or more glasses per day of milk during the past seven days							12.1	15.9	9.8	No	Yes

*Non-Hispanic.

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year										Linear Change	Quadratic Change
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
White*												
Physical Activity												
Percentage of students who were physically active for a total of at least 60 minutes per day on five or more of the past seven days								51.7	47.7		No	
Percentage of students who watched three or more hours per day of TV on an average school day								27.6	24.6		No	
Percentage of students who attended physical education (PE) classes on one or more days in an average week when they were in school							47.6	39.5	45.3		No	No
Percentage of students who attended physical education (PE) classes daily in an average week when they were in school							36.6	31.8	36.0		No	No

*Non-Hispanic.

2007 YOUTH RISK BEHAVIOR SURVEY RESULTS

North Carolina (Region 3) High School Survey
Trend Analysis Report

Question	Prevalence Survey Year										Linear Change	Quadratic Change
	1991	1993	1995	1997	1999	2001	2003	2005	2007			
White*												
Other												
Percentage of students who had ever been told by a doctor or nurse that they had asthma								19.5	19.9		No	

*Non-Hispanic.